arXiv:0907.2876v2 [math.DS] 1 Sep 2009

Bratteli-Vershik adic representations of some one-sided substitution subshifts

Reem Yassawi

Abstract

We study one-sided substitution subshifts, and how they can be respresented using Bratteli-Vershik systems. In particular we focus on minimal recognizable substitutions such that the generated one-sided substitution subshift contains only one non-shift-invertible element (branch point), and we call these substitutions *quasi-invertible*. We characterise these substitutions, and show that if the substitution is left proper, then the subshift is equal to another substitution subshift where the branch point is the substitution fixed point. We use these results to prove that any quasi-invertible substitution subshift has either a Bratteli-Vershik representation, or a 'pinched' such representation.

1. *Introduction*

The realization of any minimal homeomorphism T of a Cantor space X, as a proper Bratteli-Vershik, or adic, system, was shown in [**HPS92**]. Since then researchers have characterized adic systems with specified properties. For example, stationary proper adic systems were shown to be the union of the class of (two-sided) primitive aperiodic substitution subshifts, and the class of stationary odometers in [**VL92**], [**For97**] and [**DHS99**], and proper adic systems whose Bratteli diagram have the equal path number property were shown to be Toeplitz systems in [**GJ00**]. The adic systems realising interval exchange maps, linearly recurrent systems, and minimal substitution systems, are also described in [**GJ02**], [**CDHM03**] and [**BKM09**] respectively.

If T is only assumed to be continuous, and minimal, some constraints on the Bratteli diagrams which generate the adic representation have to be imposed. For example, while Bratteli diagrams which are no longer proper have to be considered, minimality of T implies that there have to be at least as many maximal elements in $X_{\mathcal{B}}$ as minimal elements.

In this article we address the question of how the results in [**DHS99**] can be generalized to representing one-sided (non-σ-invertible) substitution subshifts. We deal with the special case where the substitution subshift $(X_{\boldsymbol{\tau}}, \sigma)$ has a unique non-invertible point, and call these subshifts *quasi-invertible*. Subshifts with this property have been studied: Sturmian sequences, and more generally, *Arnoux-Rauzy sequences* ([**AR91**]) generate quasi-invertible subshifts - the condition (*) in [**AR91**] ensures this, and in many cases these sequences are fixed points of substitutions (see [**CMPS93**] for a characterization of when Sturmian subshifts are substitution subshifts; if the 'S-adic' expansion of a sequence in [**AR91**] is periodic, then it is the fixed point of a substitution, for example the tribonacci substitution). All these subshifts have nice geometric representations as toral rotations, or interval exchange maps. Many (but not all) Pisot substitutions that have been studied in the literature are quasi-invertible. It would be interesting to study whether quasi-invertibility imposes any

---

2000 *Mathematics Subject Classification* 37B10.

This research was partially supported by an NSERC grant.

spectral constraints. While not all have discrete spectrum (the Chacon substitution is quasi-invertible - see Example 5 in Section 5), other spectral properties may be constrained.

As invertible substitution subshifts were considered in [**DHS99**], bilateral recognizability was a sufficient assumption, and this is a property that all primitive aperiodic substitutions enjoy. Here we have to restrict ourselves to unilaterally recognizable substitutions. Thus we do not realize adic systems that arise from a non-recognizable substitution as substitution subshifts. As we want to obtain adic representations of some topological rank one systems arising from a substitution, we work with minimal substitutions. We also assume that all powers of the substitution $\boldsymbol{\tau}$ are injective. We'll call substitutions which satisfy these properties *reasonable*. We represent these systems with a stationary Bratteli diagram which has a unique maximal element. We do this by generalising the techniques in [**DHS99**], so that a reasonable quasi-invertible substitution subshift is conjugate either to an adic system, or a 'pinched' adic system.

In Section 2 we give definitions, and some background of required results for substitution subshifts. In Section 3, we characterize quasi-invertible substitutions (Lemma 9 and the remarks following it), and show that any left proper quasi-invertible substitution subshift is equal to one where the branch point is fixed by the substitution (Corollary 11. We believe that this result is still true for non-left proper subshifts -see Example (3) in Section 5). In Section 4, we set up background and notation for adic systems, and use the results of Section 3 to obtain our representation Theorem 14 for left proper quasi-invertible substitutions. We extend this result to arbitrary quasi-invertible substitutions (Theorem 19) in Section 5.

2. *Preliminaries*

2.1. *Notation*

Let $\mathcal{A}$ be a finite alphabet; elements in $\mathcal{A}$ will be denoted by a, b, c etc. A *word* $\boldsymbol{w}$ (denoted using boldface) from $\mathcal{A}$ is a finite concatenation of elements from $\mathcal{A}$. Let $\mathcal{A}^+$ denote the set of words (including the empty word) from $\mathcal{A}$. If $\boldsymbol{w} = w_1 \dots w_m$ and $\boldsymbol{w}^* = w_1^* \dots w_n^*$, then $\boldsymbol{w}\boldsymbol{w}^* := w_1 \dots w_m w_1^* \dots w_n^*$. Define $\boldsymbol{w}_{[i,j]} = w_i w_{i+1} \dots w_j$. If $\boldsymbol{w} = w_1 \dots w_m$, the *length* of $\boldsymbol{w}$, written $|\boldsymbol{w}|$, is m. Given words $\boldsymbol{w}$, $\boldsymbol{w}^*$, say that $\boldsymbol{w}$ *occurs* in $\boldsymbol{w}^*$, or $\boldsymbol{w}$ *is a subword of* $\boldsymbol{w}^*$, if $\boldsymbol{w} = {\boldsymbol{w}^*}_{[i,i+|\boldsymbol{w}|-1]}$ for some i. If $|\boldsymbol{w}| = m$, define the *k-shift* $\sigma^k(\boldsymbol{w}) = \boldsymbol{w}_{[k+1,m]}$. $\boldsymbol{w}^*$ is a *prefix* of $\boldsymbol{w}$ if $\boldsymbol{w}_{[1,|\boldsymbol{w}^*|]} = \boldsymbol{w}^*$, similarly $\boldsymbol{w}^*$ is a *suffix* of $\boldsymbol{w}$ if $\boldsymbol{w}_{[|\boldsymbol{w}|-|\boldsymbol{w}^*|+1,|\boldsymbol{w}|]} = \boldsymbol{w}^*$. When we speak of 'the' prefix (suffix) of a word $\boldsymbol{w}$, we will mean the first (last) letter of $\boldsymbol{w}$.

Let $\mathbb{N} := \{0, 1, \dots\}$. If $\mathbb{M} = \mathbb{Z}$ or $\mathbb{N}$, then the space of all $\mathbb{M}$-indexed sequences from $\mathcal{A}$ is written as $\mathcal{A}^{\mathbb{M}}$, and a configuration $\boldsymbol{x} \in \mathcal{A}^{\mathbb{M}}$ is written $\boldsymbol{x} = \{x_m\}_{m \in \mathbb{M}}$. Let $\mathcal{A}$ be endowed with the discrete topology and $\mathcal{A}^{\mathbb{M}}$ with the product topology (this topology is also generated by the Hamming metric), then $\mathcal{A}^{\mathbb{M}}$ is a *Cantor space*: a zero-dimensional perfect compact metric space. If $b \in \mathcal{A}$ and $j \in \mathbb{M}$, the clopen sets $\{\boldsymbol{x} : x_j = b\}$ form a countable basis for the topology on $\mathcal{A}^{\mathbb{M}}$. The (left) *shift map* $\sigma : \mathcal{A}^{\mathbb{M}} \to \mathcal{A}^{\mathbb{M}}$ is the map defined as $(\sigma(\boldsymbol{x}))_m = x_{m+1}$. X is a *subshift of* $(\mathcal{A}^{\mathbb{M}}, \sigma)$ if it is a closed σ-invariant subset of $\mathcal{A}^{\mathbb{M}}$. A *language* is a non-empty subset of $\mathcal{A}^+$. If $\boldsymbol{x} \in \mathcal{A}^{\mathbb{M}}$, the word $\boldsymbol{w}$ *occurs in* $\boldsymbol{x}$, or *is a word in* $\boldsymbol{x}$ if there is some $m \in \mathbb{M}$ such that $x_{m+1} \dots x_{m+|\boldsymbol{w}|} = \boldsymbol{w}$. Let $\mathcal{L}(\boldsymbol{x})$, the *language of* $\boldsymbol{x}$ be the collection of words occurring in $\boldsymbol{x}$. If $\mathcal{L}$ is a collection of words closed under the operation of taking subwords, then the set $X_{\mathbb{M}}(\mathcal{L})$ associated with $\mathcal{L}$ is the set of all sequences in $\mathcal{A}^{\mathbb{M}}$ all of whose words belong to $\mathcal{L}$. If $\mathcal{L} = \mathcal{L}(\boldsymbol{x})$, we write $X_{\mathbb{M}}(\boldsymbol{x})$, dropping the '$\mathbb{M}$' when the context makes it clear, or when a statement is true either way.

2.2. *Substitutions*

A *substitution* is a map $\boldsymbol{\tau} : \mathcal{A} \to \mathcal{A}^+$. We extend $\boldsymbol{\tau}$ to a map $\boldsymbol{\tau} : \mathcal{A}^+ \to \mathcal{A}^+$ by concatenation: if $\boldsymbol{a} = a_1 \dots a_k$, then $\boldsymbol{\tau}(\mathbf{a}) := \boldsymbol{\tau}(a_1) \dots \boldsymbol{\tau}(a_k)$. In this way iteration $\boldsymbol{\tau}^n$ is well defined. The substitution $\boldsymbol{\tau}$ is extended to a map $\boldsymbol{\tau} : \mathcal{A}^{\mathbb{N}} \to \mathcal{A}^{\mathbb{N}}$ defined by $\boldsymbol{\tau}(\boldsymbol{x}) := \boldsymbol{\tau}(x_0)\,\boldsymbol{\tau}(x_1)\dots$, and also $\boldsymbol{\tau} : \mathcal{A}^{\mathbb{Z}} \to \mathcal{A}^{\mathbb{Z}}$ defined by $\boldsymbol{\tau}(\boldsymbol{x}) := \dots \boldsymbol{\tau}(x_{-1}) \cdot \boldsymbol{\tau}(x_0)\,\boldsymbol{\tau}(x_1) \dots$. We say $\boldsymbol{\tau}$ is *left (right) proper* if there exists l in $\mathcal{A}$ such that l is a prefix (suffix) of $\boldsymbol{\tau}(a)$ for each $a \in \mathcal{A}$. If $\boldsymbol{\tau}$ is both left and right proper, it is called *proper*. We say $\boldsymbol{\tau}$ *suffix-permutative* if the set of suffixes of $\{\boldsymbol{\tau}(a)\}_{a\in\mathcal{A}}$ is a permutation of $\mathcal{A}$. Also $\boldsymbol{\tau}$ is *injective* if $\boldsymbol{\tau}(a) \neq \boldsymbol{\tau}(b)$ whenever $a \neq b$.

We say $\boldsymbol{\tau}$ is *primitive* if there exists a positive integer k such that for any $a \in \mathcal{A}$, all letters of $\mathcal{A}$ appear in $\boldsymbol{\tau}^k(a)$ (this requires that for some letter a, $l_a > 1$). If $\mathcal{L}_{\boldsymbol{\tau}}$ is the language generated by the words $\{\boldsymbol{\tau}^n(a) : n \in \mathbb{N}^+,\ a \in \mathcal{A}\}$, then $X_{\mathbb{M}}(\boldsymbol{\tau}) := X_{\mathbb{M}}(\mathcal{L}_{\boldsymbol{\tau}})$ is closed and σ-invariant, so that $(X_{\mathbb{M}}(\boldsymbol{\tau}), \sigma)$ is a (substitution) subshift, with the subspace topology. Henceforth we will be working mainly with one sided substitution subshifts, and unless otherwise indicated, will assume this, and write $(X_{\boldsymbol{\tau}}, \sigma)$ instead of $(X_{\mathbb{N}}(\boldsymbol{\tau}), \sigma)$. A *fixed point* of $\boldsymbol{\tau}$ is a sequence $\boldsymbol{u} \in \mathcal{A}^{\mathbb{N}}$ such that $\boldsymbol{\tau}(\boldsymbol{u}) = \boldsymbol{u}$. If $\boldsymbol{u} \in \mathcal{A}^{\mathbb{N}}$ is a fixed point for $\boldsymbol{\tau}$, then $\boldsymbol{\tau}(u_0)$ starts with u_0. Conversely, if there exists a letter l such that l is a prefix of $\boldsymbol{\tau}(l)$, then $\boldsymbol{u} := \lim_{n\to\infty} \boldsymbol{\tau}^n(l)$ is the unique fixed point satisfying $u_0 = l$. If $\boldsymbol{\tau}$ is primitive, then the subshift generated by $\boldsymbol{u}$, i.e. $\overline{\{\sigma^n(\boldsymbol{u})\}}$, equals $X_{\boldsymbol{\tau}}$. Using the pigeonhole principle, there exists $n \geq 1$ such that $\boldsymbol{\tau}^n$ has at least one fixed point. Though we do not assume that $\boldsymbol{\tau}$ is primitive, we will assume that $\boldsymbol{\tau}$ has a fixed point $\boldsymbol{u}$ with $\mathcal{L}_{\boldsymbol{\tau}} = \mathcal{L}(\mathbf{u})$, and $|\boldsymbol{\tau}^n(u_0)| \to \infty$ as $n \to \infty$. When such a fixed point exists, which we henceforth assume, we will call it a *generating* fixed point. Note that if $\boldsymbol{\tau}$ is left proper and has a generating fixed point, then it is primitive.

The continuous mapping $T : X \to X$ of the compact metric space X is *minimal* if for each x in X, $\{T^n(x) : n \geq 0\}$ is dense in X. Primitive substitutions are minimal. A sequence $\boldsymbol{x} \in \mathcal{A}^{\mathbb{M}}$ is *almost periodic* if for any neighbourhood V of $\boldsymbol{x}$, the set $\{n : \sigma^n(\boldsymbol{x}) \in V\}$ has bounded gaps in $\mathbb{M}$. If $\boldsymbol{u}$ is almost periodic, then $(X(\boldsymbol{u}), \sigma)$ is minimal. (see [**Que87**], Theorem 4.12). In what follows we will assume that our substitution $\boldsymbol{\tau}$ has a generating fixed point $\boldsymbol{u}$, and that the resulting subshift (X, σ) is minimal. We call such substitutions *minimal*. Minimal substitutions enjoy many established properties that primitive substitutions do, and minor modifications of proofs of facts for primitive substitutions can be made to ensure that the same results hold for minimal substitutions. In particular unilateral recognizability, (see below) is defined for primitive substitutions. The necessary and sufficient condition that exists for a primitive substitution to be unilaterally recognizable can be modified with the following two lemmas (which appear in [**Mos92**] for primitive substitutions):

LEMMA 1. *Fix a natural number l. If $\boldsymbol{w}$ occurs in $\boldsymbol{u}$, $|\boldsymbol{w}| = l$, and $\boldsymbol{\tau}^p(\boldsymbol{w}) = B_p \boldsymbol{\tau}^p(\boldsymbol{w}^*) B'_p$, with $\boldsymbol{w}^*$ occurring in $\boldsymbol{u}$ and $|\boldsymbol{w}^*| = l^*$, then there is C such that $l^* \leq C\,l$, where C is independent of p.*

Proof. Suppose not. Then for all N there is a p, a $\boldsymbol{w}^* \in \mathcal{L}_{\boldsymbol{\tau}}$ such that $\boldsymbol{\tau}^p(\boldsymbol{w}) = B_p \boldsymbol{\tau}^p(\boldsymbol{w}^*) B'_p$ with $|\boldsymbol{w}^*| = l^*$ and $l^* > Nl$. If $\boldsymbol{w} = w_1 \dots w_l$ and $\boldsymbol{w}^* = w^*_1 \dots w^*_{l^*}$, then $\left|\prod_{i=1}^{l} \boldsymbol{\tau}^p(w_i)\right| \geq \left|\prod_{i=1}^{l^*} \boldsymbol{\tau}^p(w^*_i)\right|$. Since $\boldsymbol{\tau}$ is minimal, $\boldsymbol{u}$ is almost periodic, which means that all letters occur in $\boldsymbol{u}$ with bounded periodicity. Hence if for $\alpha \in \mathcal{A}$ we let $n(\alpha)$ denote the number of occurrences of α in $\boldsymbol{w}$, then there are more than $n(\alpha)$ occurrences of α in $\boldsymbol{w}^*$ for each α, if N is large enough. This contradicts the inequality above. ■

Lemma 2. *Fix a natural k. The sequence*

$$L_p := \frac{\max\{|\boldsymbol{\tau}^p(\boldsymbol{w})| : \boldsymbol{w} \in \mathcal{L}_{\boldsymbol{u}}, |\boldsymbol{w}| = k\}}{\min\{|\boldsymbol{\tau}^{p-1}(a)| : a \in \mathcal{A} \text{ and } \lim_{n\to\infty} \boldsymbol{\tau}^n(a) = \infty\}}$$

is bounded.

Proof. Let $\mathcal{A}_\infty := \{a \in \mathcal{A} : \lim_{n\to\infty} |\boldsymbol{\tau}^n(a)| = \infty\} = \{\alpha_1, \dots \alpha_{|\mathcal{A}_\infty|}\}$. It is sufficient to show that for some C, $|\boldsymbol{\tau}^p(\alpha)| \leq C|\boldsymbol{\tau}^p(\beta)|$ for all natural p and all $\alpha, \beta \in \mathcal{A}_\infty$. If M is the $|\mathcal{A}_\infty|$-sized square matrix whose i, j-indexed entry is the number of occurrences of α_j in $\boldsymbol{\tau}(\alpha_i)$, then wlog, M can be assumed to have all positive entries, and the Perron-Frobenius theorem [**Sen06**] implies that $|\{\beta \in \boldsymbol{\tau}^p(\alpha) : \beta \in \mathcal{A}_\infty\}| = C_\alpha \lambda^p + o(\lambda^p)$, where C_α is strictly positive and λ is the dominant eigenvalue of M. Thus there is a C such that

$$|\{\gamma \in \boldsymbol{\tau}^p(\alpha) : \gamma \in \mathcal{A}_\infty\}| \leq C|\{\gamma \in \boldsymbol{\tau}^p(\beta) : \gamma \in \mathcal{A}_\infty\}|$$

whenever $\alpha, \beta \in \mathcal{A}_\infty$. The result follows from the almost periodicity of $\boldsymbol{u}$. ■

Unilateral recognizability was a condition introduced in [**Hos86**] and [**Que87**] which ensured that one could find clopen generating partitions for $(X_{\boldsymbol{\tau}}, \sigma)$. Specifically, the set of *(1)-cuttings* of $\boldsymbol{u}$ are

$$E := \{0\} \cup \{|\boldsymbol{\tau}(\boldsymbol{u}_{[0,p-1]})| : p > 0\}.$$

We then say that $\boldsymbol{\tau}$ is *(unilaterally) recognizable* if there exists some L so that if $\boldsymbol{u}_{[i,i+L-1]} = \boldsymbol{u}_{[j,j+L-1]}$ and $i \in E$, then $j \in E$. A word $\boldsymbol{w}$ occurring at $\mathbf{u}_i$ and $\boldsymbol{u}_j$ has the *same 1-cutting* at i and j, with $i < j$, if $E \cap \{i, \dots i + |\boldsymbol{w}|\} = j - i + E \cap \{j, \dots j + |\boldsymbol{w}|\}$.

In Theorem 3.1, [**Mos92**], primitive recognizable substitutions are characterized, and Lemmas 1 and 2 can be used to extend this theorem to minimal recognizable substitutions:

Theorem 3. *Let $\boldsymbol{\tau}$ be minimal with generating fixed point $\boldsymbol{u}$. $\boldsymbol{\tau}$ is not recognizable if and only if for each L, there exists a word $\boldsymbol{w}$ of length L, and two elements a, $b \in \mathcal{A}$ such that*

- *$\boldsymbol{\tau}(b)$ is a proper suffix of $\boldsymbol{\tau}(a)$.*
- *The words $\boldsymbol{\tau}(a)\boldsymbol{w}$ and $\boldsymbol{\tau}(b)\boldsymbol{w}$ appear in $\boldsymbol{u}$ with the same 1-cutting of $\boldsymbol{w}$.* ■

In particular, suffix permutative substitutions are recognizable. It is straightforward to show that if $\boldsymbol{\tau}$ is recognizable, and injective, then $\boldsymbol{\tau}^n$ is recognizable for each n.

In Lemmas 2 and 3, [**Hos86**], the following is shown for primitive recognizable substitutions. Note that $\boldsymbol{\tau}(X) \subset X$.

Proposition 4.
Let $\boldsymbol{\tau}$ be a primitive, recognizable substitution with generating fixed point $\boldsymbol{u}$.

(i) *$\boldsymbol{y} \in \boldsymbol{\tau}(X)$ if and only if $\sigma^{n_i}\boldsymbol{u} \to \boldsymbol{y}$ where $n_i \in E$ for all large i.*
(ii) *$\boldsymbol{\tau}(X)$ is clopen in X.*
(iii) *$\sigma^p(\boldsymbol{\tau}(x)) \in \boldsymbol{\tau}(X)$ if and only if $p = |\boldsymbol{\tau}(\boldsymbol{x}_{[0,r]})|$ for some r so that $\sigma^p(\boldsymbol{\tau}(x)) = \boldsymbol{\tau}(\sigma^{r+1}(\boldsymbol{x}))$.*

If $\boldsymbol{\tau}$ is also injective, then

(iv) *$\boldsymbol{\tau} : X \to \boldsymbol{\tau}(X)$ is a homeomorphism.* ■

We remark here that the previous proposition also implies that for each $n \geq 1$, $\boldsymbol{\tau}^n : X \to \boldsymbol{\tau}^n(X)$ is also a homeomorphism, though as a substitution it is not necessarily injective.

Let us call substitutions which are recognizable, minimal, and all of whose powers are injective *reasonable*. In some of what follows (excepting Corollary 12, and all results that require it), we need only assume that $\boldsymbol{\tau}$ is injective, but since our final results assume that all powers of $\boldsymbol{\tau}$ are injective, we will make this blanket assumption everywhere.

Corollary 5. *Suppose that $\boldsymbol{\tau}$ is reasonable. Every $\boldsymbol{y} \in \boldsymbol{\tau}(X)$ can be written in a unique way as $\mathbf{y} = \boldsymbol{\tau}(\boldsymbol{x})$ with $\boldsymbol{x} \in X$. Every $\boldsymbol{y} \in X\backslash\boldsymbol{\tau}(X)$ can be written as $\boldsymbol{y} = \sigma^k\boldsymbol{\tau}(\boldsymbol{x})$ where $\sigma(\boldsymbol{x})$ is unique, and $0 < k < |\boldsymbol{\tau}(x_0)|$.*

Proof. The first assertion follows immediately from Proposition 4. If $\boldsymbol{y} \in X\backslash\boldsymbol{\tau}(X)$ then by Part 1 of Proposition 4, there is a sequence $\{n_i\}$ of integers not in E, with $\sigma^{n_i}(\boldsymbol{u}) \to \boldsymbol{y}$. Now each $n_i = k_i + r_i$ where $k_i \in E$, and the positive r_i is strictly less than the successor of k_i in E. Since E is almost periodic, by dropping to a subsequence if necessary, we assume that there is some fixed positive r such that each $n_i = k_i + r$. Now we can assume that $\sigma^{k_i}(\boldsymbol{u})$ converges to some $\boldsymbol{x} \in \boldsymbol{\tau}(X)$. So $\boldsymbol{y} = \sigma^r\boldsymbol{x}$.

Now suppose that $\boldsymbol{y} = \sigma^r\boldsymbol{x} = \sigma^{r'}\boldsymbol{x}'$, where if $\boldsymbol{x} = x_0x_1\ldots$ then $r < |\boldsymbol{\tau}(x_0)|$, and similarly for r'. Let $l = |\boldsymbol{\tau}(x_0)| - r$, and define l' similarly.

If $l = l'$, then $\sigma^{r+l}\boldsymbol{\tau}(\boldsymbol{x}) = \boldsymbol{\tau}(\sigma\boldsymbol{x}) \in \boldsymbol{\tau}(X)$ and similarly $\sigma^{r'+l}\boldsymbol{\tau}(x') = \boldsymbol{\tau}(\sigma\boldsymbol{x}') \in \boldsymbol{\tau}(X)$, so $\boldsymbol{\tau}(\sigma\boldsymbol{x}) = \boldsymbol{\tau}(\sigma\boldsymbol{x}')$. Thus $\sigma\boldsymbol{x} = \sigma\boldsymbol{x}'$.

Finally we will show that $l \neq l'$ leads to a contradiction. Suppose $l < l'$. Then $\sigma^{r+l}\boldsymbol{\tau}(\boldsymbol{x}) = \boldsymbol{\tau}(\sigma\boldsymbol{x}) \in \boldsymbol{\tau}(X)$, and so $\sigma^{r'+l}\boldsymbol{\tau}(\boldsymbol{x}') = \sigma^{r+l}\boldsymbol{\tau}(\boldsymbol{x}) \in \boldsymbol{\tau}(X)$. Thus by Part 3 of Proposition 4, $r' + l = |\boldsymbol{\tau}(\boldsymbol{x}'_{[0,j]})|$ for some j. This contradicts $r' + l < r' + l' = |\boldsymbol{\tau}(x'_0)|$. ■

3. *Quasi-invertible substitutions*

We investigate the number of σ-preimages that a $\boldsymbol{\tau}$-fixed point $\boldsymbol{u}$ can have. Say that a_1 is *part of a suffix cycle* $a_1, a_2, \ldots a_m$ if $\boldsymbol{\tau}(a_i)$ has a_{i+1} as a suffix for $1 \le i < m$, and $\boldsymbol{\tau}(a_m)$ has a_1 as a suffix.

Proposition 6. *Suppose that $\boldsymbol{\tau}$ is a reasonable substitution with generating fixed point $\boldsymbol{u} = u_0u_1u_2\ldots$. Then $a\boldsymbol{u} \in X_{\boldsymbol{\tau}}$ if and only if a is part of a suffix cycle $a_1, a_2, \ldots a_m$, and $a_iu_0 \in \mathcal{L}_{\boldsymbol{\tau}}$ for some $i \in \{1, \ldots m\}$.*

Proof. If for some i, $a_iu_0 \in \mathcal{L}_{\boldsymbol{\tau}}$, then since $\boldsymbol{u}$ is generating, there exists an n such that $a_iu_0 \in \boldsymbol{\tau}^n(u_0)$, so $a_{i+1}\,\boldsymbol{\tau}(u_0) \in \boldsymbol{\tau}^{n+1}(u_0)$, and $a_i\boldsymbol{\tau}^{nm}(u_0)$ for each $n \ge 1$. It follows that $a_i\boldsymbol{u} \in X_{\boldsymbol{\tau}}$, and that this is true for each $a_j \in \{a_1, \ldots a_m\}$.

Conversely, suppose that $a\boldsymbol{u} \in X$. We claim first that a is the suffix of some substitution word. Now by Corollary 5, $a\boldsymbol{u} = \boldsymbol{w}\boldsymbol{\tau}(\sigma\boldsymbol{x})$ where $\boldsymbol{w}$ is the suffix of some substitution word $\boldsymbol{\tau}(x_0)$, and $\sigma\boldsymbol{x}$ is unique. Thus $\boldsymbol{\tau}(\boldsymbol{u}) = \boldsymbol{u} = \sigma(\boldsymbol{w})\boldsymbol{\tau}(\sigma\boldsymbol{x})$, and by Corollary 5 again, $\sigma(\boldsymbol{w}) = \boldsymbol{\tau}(u_0 \ldots u_r)$ for some $r \ge 0$, so that a is the suffix of a substitution word.

Since (X, σ) is minimal, σ is onto, which means that for each n there are words $\boldsymbol{w}_n$ of length n such that $\boldsymbol{w}_na\boldsymbol{u} \in X$. Given N, choose n so that $n > |\boldsymbol{\tau}^N(\alpha)|$ for all α. Thus $\boldsymbol{x} = \boldsymbol{w}_na\boldsymbol{u} = \boldsymbol{w}'\boldsymbol{\tau}^N(\sigma(\boldsymbol{y}))$ where $\boldsymbol{w}'$ is the suffix of $\boldsymbol{\tau}^N(y_0)$, and Corollary 5 implies that for some $p > 0$, $\sigma^{p+1}(\boldsymbol{y}) = \boldsymbol{u}$ and $\boldsymbol{w}_na = \boldsymbol{w}'\boldsymbol{\tau}^N(\boldsymbol{y}_{[1,p]})$, so that a is the suffix of a $\boldsymbol{\tau}^N$ word. Taking a subsequence, we conclude that there is some b such that a is the suffix of $\boldsymbol{\tau}^N(b)$ for infinitely many N's. Since the sequence of suffixes of $\boldsymbol{\tau}^N(b)$ is eventually periodic, this implies that a is part of a suffix cycle. ■

It follows that if $\boldsymbol{\tau}$ is left proper and suffix permutative, then $\boldsymbol{u}$ has $|\mathcal{A}|$ σ-preimages, and if $\boldsymbol{\tau}$ is left and right proper, then $\boldsymbol{u}$ has one σ-preimage.

Let $\mathcal{W}_n := \{\boldsymbol{w} : \boldsymbol{w} = \sigma^k(\boldsymbol{\tau}^n(a)) : a \in \mathcal{A},\ 0 \leq k < |\boldsymbol{\tau}^n(a)|\}$. If $\boldsymbol{w} \in \mathcal{W}_n$, let $r(\boldsymbol{w})$ be the number of distinct representations $\boldsymbol{w}$ has as $\boldsymbol{w} = \sigma^k(\boldsymbol{\tau}^n(a))$, as k and a vary, and let $\sigma^{-1}\boldsymbol{w} = \{\alpha\boldsymbol{w} : \alpha\boldsymbol{w} \in \mathcal{W}_n\}$.

Proposition 7. *Let $\boldsymbol{\tau}$ be reasonable, with a generating fixed point $\boldsymbol{u}$. Suppose that $\boldsymbol{y} \neq \boldsymbol{u}$, $\boldsymbol{y} \in [\boldsymbol{w}]$ where $\boldsymbol{w} = \sigma^k\boldsymbol{\tau}^n(a)$, and $\boldsymbol{w}' := \sigma^{k-1}\boldsymbol{\tau}^n(a)$. Then $\boldsymbol{y}$ has more than one σ preimage only if $r(\boldsymbol{w}') < r(\boldsymbol{w})$.*

Conversely, suppose that there exists a sequence of words $\boldsymbol{w}_n \in \mathcal{W}_n$ with $|\boldsymbol{w}_n| \to \infty$, with $\cap_n[\boldsymbol{w}_n] = \boldsymbol{y}$. If there exists a sequence a_n such that $a_n\boldsymbol{w}_n \in \mathcal{W}_n$ and $r(a_n\boldsymbol{w}_n) < r(\boldsymbol{w}_n)$ for each n, then $\boldsymbol{y}$ has more than one σ-preimage.

Proof. The only way that $\boldsymbol{y}$ can have more than one σ-preimage is if $\boldsymbol{y}$ has at least two representations $\boldsymbol{y} = \sigma^k\boldsymbol{\tau}^n(a)\boldsymbol{x} = \sigma^{k'}\boldsymbol{\tau}^n(b)\boldsymbol{x}$ with $\sigma^k\boldsymbol{\tau}^n(a) = \sigma^{k'}\boldsymbol{\tau}^n(b) = \boldsymbol{w}$ but $\sigma^{k-1}\boldsymbol{\tau}^n(a) \neq \sigma^{k'-1}\boldsymbol{\tau}^n(b)$.

Conversely, note that if $r(\alpha\boldsymbol{w}) < r(\boldsymbol{w})$, for some $\alpha\,\boldsymbol{w} \in \sigma^{-1}\boldsymbol{w}$, this means that there is more than one word in $\sigma^{-1}\boldsymbol{w}$. So for each n there are two distinct letters α_n and β_n so that $\alpha_n\boldsymbol{w}_n$ and $\beta_n\boldsymbol{w}_n$are both in $\mathcal{W}_n$. Passing down to a subsequence if necessary, we can assume that there are distinct α and β so that $\alpha\boldsymbol{w}_n$ and $\beta\boldsymbol{w}_n$ are elements of $\mathcal{W}_n$, hence $\mathcal{L}_{\boldsymbol{\tau}}$. Thus $\alpha\boldsymbol{y}$ and $\beta\boldsymbol{y}$ are both elements of X, so that $\boldsymbol{y}$ has more than one σ-preimage. ■

Corollary 8. *Suppose that $\boldsymbol{\tau}$ is a reasonable substitution with generating fixed point $\boldsymbol{u}$. Then*

- (i) *If $\boldsymbol{\tau}$ is suffix permutative then only $\boldsymbol{u}$ in X has more than one σ-preimage.*
- (ii) *If $\boldsymbol{\tau}$ is left proper, and not suffix permutative, such that whenever $\boldsymbol{\tau}(a)$, $\boldsymbol{\tau}(b)$ share a proper common suffix, then $\boldsymbol{\tau}(b)$ is a proper suffix of $\boldsymbol{\tau}(a)$ (or vice versa), then only $\boldsymbol{u}$ has more than one σ-preimage.*
- (iii) *Suppose $\boldsymbol{\tau}$ is left proper, not suffix permutative, and there are a, b such that $\boldsymbol{\tau}(a)$, $\boldsymbol{\tau}(b)$ share a common proper suffix $\mathbf{s}$. If $\boldsymbol{\tau}^n(a)$, $\boldsymbol{\tau}^n(b)$ share the common suffix $\mathbf{s}_n$, then $\boldsymbol{y} := \lim_{n\to\infty}\mathbf{s}_n$, if it exists, has more than one σ-preimage. Also $\boldsymbol{y} = \boldsymbol{u}$ if and only if $\mathbf{s}_n \in \boldsymbol{\tau}^{k_n}(X)$ for some $k_n \to \infty$.*

Proof. Suppose that $\boldsymbol{\tau}$ is suffix permutative. Then no two $\boldsymbol{\tau}^n$-words share a common suffix. So $r(\boldsymbol{w}) = 1$ for all $\boldsymbol{w} \in \mathcal{W}_n$ and all n. By Proposition 7, other than $\boldsymbol{u}$, there are no other points with more than one σ-preimage.

Suppose that $\boldsymbol{\tau}$ is not suffix permutative. Proposition 7 tells us that the only candidates for a point $\boldsymbol{y}$ with more than one σ-preimage are those where $\boldsymbol{y} = \lim_n \mathbf{s}_n$ with $\mathbf{s}_n$ being the suffix of more than one $\boldsymbol{\tau}_n$ word. For this limit to exist, the lengths of the words $\mathbf{s}_n$ must tend to ∞.

If any two substitution words $\boldsymbol{\tau}(a)$, $\boldsymbol{\tau}(b)$ share a common suffix only when $\boldsymbol{\tau}(b)$ is a (proper) suffix of $\boldsymbol{\tau}(a)$, then $\boldsymbol{\tau}^n(b)$ is a suffix of $\boldsymbol{\tau}^n(a)$ for every n. Suppose $|\boldsymbol{\tau}^n(\mathbf{b})| \to \infty$, so that $\boldsymbol{y} = \lim_{n\to\infty}\boldsymbol{\tau}^n(b)$ is a candidate for a point with multiple σ-preimages. Then $\boldsymbol{y} \in \boldsymbol{\tau}^n(X)$ for each n, so $\boldsymbol{y} = \boldsymbol{u}$.

If a non suffix-permutative $\boldsymbol{\tau}$ is such that no substitution word is the (proper) suffix of another, then there are distinct a and b such that for large n $\boldsymbol{\tau}^n(a)$ and $\boldsymbol{\tau}^n(b)$ share a common suffix $\mathbf{s}_n$, and assuming their lengths tend to ∞, these words satisfy the conditions of Proposition 7. Thus there exists a point $\boldsymbol{y}$ with more than one σ-preimage. Now $\boldsymbol{y} = \boldsymbol{u}$ if and only if $\boldsymbol{y} \in \cap \boldsymbol{\tau}^n(X)$, from which the result follows. ■

Now that we have partial conditions under which a reasonable substitution $\boldsymbol{\tau}$ has a unique, non-fixed non-σ-invertible point y, we recast $(X_{\boldsymbol{\tau}}, \sigma)$ as a substitution system for which $\boldsymbol{y}$ is fixed. We need to introduce some notation. If $\boldsymbol{\tau}$ is a right proper, reasonable substitution on the alphabet $\mathcal{A} = \{a_1, \dots, a_n\}$, let $\mathbf{s}_1$ be the maximal proper common suffix of $\{\boldsymbol{\tau}(a)\}_{a \in \mathcal{A}}$, so that for each i in $\mathcal{A}$, $\boldsymbol{\tau}(a_i) = \boldsymbol{a}_i x_i \mathbf{s}_1$, where the set $\{x_1, \dots x_n\}$ has at least two distinct elements. Call the set $A_1 := \{x_i\}_{i=1}^n$ the set of $\mathbf{s}_1$-*predecessors* in $\{\boldsymbol{\tau}(a)\}_{a \in A}$. If $\{\boldsymbol{\tau}(x_i)\}_{x_i \in A_1}$ has $\mathbf{e}_2\mathbf{s}_1$ as a maximal proper common suffix ($\mathbf{e}_2$ could be empty), let A_2 be the set of $\mathbf{e}_2\mathbf{s}_1$-predecessors in $\{\boldsymbol{\tau}(a)\}_{a \in A_1}$. Inductively, if A_k is the set of $\mathbf{e}_k\mathbf{s}_1$-predecessors, let $\mathbf{e}_{k+1}\mathbf{s}_1$ be the maximal proper common suffix of $\{\boldsymbol{\tau}(a)\}_{a \in A_k}$, and A_{k+1} the set of $\mathbf{e}_{k+1}\mathbf{s}_1$-predecessors in $\{\boldsymbol{\tau}(a)\}_{a \in A_k}$. The sequence A_k is eventually periodic. For, $|A_k|$ cannot increase. If $|A_k|$ stays constant as k increases, then for some k and m, $A_k = A_{k+m}$, and so $\mathbf{e}_{k+1}\mathbf{s}_1 = \mathbf{e}_{k+m+1}\mathbf{s}_1$, which in turn implies that $A_{k+1} = A_{k+m+1}$, and by induction that A_k is eventually periodic. If $|A_k|$ decreases, then repeat this argument. Thus $\{\mathbf{e}_k\mathbf{s}_1\}_{k \in \mathbb{N}}$ is also eventually periodic. More generally, if $\boldsymbol{\tau}$ is not right proper, and if $\mathcal{A}^* \subset \mathcal{A}$, we can repeat the above procedure to obtain an $\mathbf{s}_1^*$, and sequences $\{\mathbf{e}_k^*\}_{k \geq 2}$ and $\{A_k^*\}_{k \geq 1}$; as above these sequences are eventually periodic. If (Y, T) is a system where y is not T-invertible, we call y a *branch point*. Let us call systems (Y, T) with only one branch point *quasi-invertible*. If $(X_{\boldsymbol{\tau}}, \sigma)$ is quasi-invertible, and the branch point has M-σ-preimages, we'll call $\boldsymbol{\tau}$ *M-quasi-invertible*. Corollary 8 characterizes these substitutions when they are left proper. We use this notation in the following lemma:

LEMMA 9. *Let $\boldsymbol{\tau}$ be a reasonable substitution on the alphabet $\mathcal{A} = \{a_1, \dots, a_n\}$, with fixed point $\boldsymbol{u}$, and with $\mathbf{s}_1$, $\{e_k\}_{k=1}^\infty$ and $\{A_k\}_{k=1}^\infty$ as defined. Suppose that $\mathbf{e}_{k+p}\mathbf{s}_1 = \mathbf{e}_k\mathbf{s}_1$ for $k \geq M + 1$. Then*

(i) *If the $\boldsymbol{\tau}$-substitution words have a maximal non-empty proper common suffix $\mathbf{s}_1$, then for $0 \leq l \leq M + p - 1$, the $\boldsymbol{\tau}^l$-substitution words have a maximal proper common suffix $\left(\prod_{k=0}^{l-2} \boldsymbol{\tau}^k(\mathbf{e}_{l-k}\mathbf{s}_1)\right) \boldsymbol{\tau}^{l-1}(\mathbf{s}_1)$, and for $n \geq 1$, the $\boldsymbol{\tau}^{np+M-1}$-substitution words have a maximal proper common suffix $\mathbf{s}_{np+M-1} = \mathbf{P}_{np+M-1}\,\mathbf{I}_{np+M-1}$, where*

$$\mathbf{P}_{np+M-1} = \prod_{k=0}^{n-1} \boldsymbol{\tau}^{pk}\left((\mathbf{e}_{M+p}\mathbf{s}_1)\boldsymbol{\tau}(\mathbf{e}_{M+p-1}\mathbf{s}_1)\boldsymbol{\tau}^2(\mathbf{e}_{M+p-2}\mathbf{s}_1)\dots\boldsymbol{\tau}^{p-1}(\mathbf{e}_{M+1}\mathbf{s}_1)\right) \tag{3.1}$$

and

$$\mathbf{I}_{np+M-1} = \left(\prod_{k=0}^{M-3} \boldsymbol{\tau}^{np+k}(\mathbf{e}_{M-1-k}\mathbf{s}_1)\right)\boldsymbol{\tau}^{np+M-2}(\mathbf{s}_1). \tag{3.2}$$

(ii) *Suppose that $\boldsymbol{\tau}$ is left proper. Then $\boldsymbol{\tau}$ is quasi-invertible, with branch point $\boldsymbol{y} \neq \boldsymbol{u}$, if and only if for some m, the $\boldsymbol{\tau}^m$-substitution words have a maximal proper non-empty common suffix $\mathbf{s}_1$, $p = 1$, and for all subsets $\mathcal{A}^* \subset \mathcal{A}$, $\mathbf{e}_{k+1}^*\mathbf{s}_1^* = \mathbf{e}_{M+1}\mathbf{s}_1$ for all k large.*

(iii) *Suppose $\boldsymbol{\tau}$ is left proper. Then $\boldsymbol{\tau}$ is quasi-invertible, with branch point $\boldsymbol{u}$, if and only if for all subsets $\mathcal{A}^* \subset \mathcal{A}$, there is some N such that for $n > N$, the $\boldsymbol{\tau}$-substitution words of elements A_n have no proper maximal common suffix.*

Proof.

(1) The proof of Equations 3.1 and 3.2 follow from the definitions of $\mathbf{e}_k\mathbf{s}_1$, and induction.

(2) If $\boldsymbol{y}$ is the branch point and $\boldsymbol{y} \neq \boldsymbol{u}$, then $\boldsymbol{u}$ is σ-invertible, so by Proposition 6, working with a higher power of $\boldsymbol{\tau}$ if necessary, we can assume that the the $\boldsymbol{\tau}$-substitution words have a maximal common suffix $\mathbf{s}_1$, and $\mathbf{s}_1$ has to be proper, or else $(X_{\boldsymbol{\tau}}, \sigma)$ is periodic

and so invertible. If $\boldsymbol{\tau}$ is such that $M = 1$, then Equation 3.1 reduces to

$$\mathbf{s}_{np} = \mathbf{P}_{np} = \prod_{k=0}^{n-1} \boldsymbol{\tau}^{pk}\left((\mathbf{e}_{p+1}\mathbf{s}_1)\boldsymbol{\tau}(\mathbf{e}_p\mathbf{s}_1)\boldsymbol{\tau}^2(\mathbf{e}_{p-1}\mathbf{s}_1)\dots\boldsymbol{\tau}^{p-1}(\mathbf{e}_2\mathbf{s}_1)\right). \tag{3.3}$$

Here $\mathbf{s}_{(n+1)p} \subset \mathbf{s}_{np}$ for $n \geq 1$, and if the $\boldsymbol{\tau}$-substitution words start with u_0, then $u_0 \in \mathbf{s}_{np}$ for $n > 1$, so that $|\mathbf{s}_{np}| \to \infty$. Thus $\boldsymbol{y} := \lim_{n\to\infty} \mathbf{s}_{np}$ exists, and by Proposition 7 has at least 2 σ-preimages. If $p > 1$, a formula similar to Equation 3.1 for $\{\mathbf{s}_{np+k}\}_{n\geq 0}$ can be obtained for $0 < k < p$. These words start with $\mathbf{e}_k\mathbf{s}_1 \neq \mathbf{e}_p\mathbf{s}_1$, leading to another branch point. Even if $M > 1$, then the initial portion $\mathbf{P}_n$ of $\mathbf{s}_n$ still converges to a branch point $\boldsymbol{y}$, and $\boldsymbol{y}$ also equals $\lim_{n\to\infty} \mathbf{s}_n$. Finally, if $\mathcal{A}^* \subset \mathcal{A}$ is such that $e^*_{k+1}s^*_1 \neq e_{M+1}s_1$ for infinitely many k, then

$$\boldsymbol{y}^* = \lim_{n\to\infty} \mathbf{P}^*_{np} = \prod_{k=0}^{n-1} \boldsymbol{\tau}^{pk}\left((\mathbf{e}^*_{p+1}\mathbf{s}^*_1)\boldsymbol{\tau}(\mathbf{e}^*_p\mathbf{s}^*_1)\boldsymbol{\tau}^2(\mathbf{e}^*_{p-1}\mathbf{s}^*_1)\dots\boldsymbol{\tau}^{p-1}(\mathbf{e}^*_2\mathbf{s}^*_1)\right)$$

also is a branch point. Note that $\boldsymbol{y}^* = \mathbf{e}^*_{M^*+1}\mathbf{s}^*_1\boldsymbol{\tau}(\boldsymbol{y}^*)$, while $\mathbf{e}_M\mathbf{s}_1\boldsymbol{\tau}(\boldsymbol{y}) = \boldsymbol{y}$, so that $\boldsymbol{y} \neq \boldsymbol{y}^*$ (this is assuming $p = 1$; the case $p > 1$ is similar).

Conversely, if the $\boldsymbol{\tau}$-substitution words $\boldsymbol{\tau}(a_i)$ have a maximal proper non-empty common suffix $\mathbf{s}_1$, and $p = 1$, then by Part 1, the $\boldsymbol{\tau}^{n+M-1}$-substitution words have maximal proper common suffixes $\mathbf{s}_{n+M-1} = \mathbf{P}_{n+M-1}\mathbf{I}_{n+M-1}$ for each n, and, as above, $\mathbf{P}_{n+M-1} \to \boldsymbol{y}$ as $n \to \infty$, with $|\sigma^{-1}(\boldsymbol{y})| > 1$. The condition on subsets $\mathcal{A}^*$ of $\mathcal{A}$ prevents there being any other branch points, and, since $\mathbf{e}_{M+1}\mathbf{s}_1\boldsymbol{\tau}(\boldsymbol{y}) = \boldsymbol{y}$ and $\boldsymbol{\tau}(\boldsymbol{u}) = \boldsymbol{u}$, $\boldsymbol{y} \neq \boldsymbol{u}$.

(3) If $\boldsymbol{u}$ is the branch point, then by Proposition 6,there are at least two distinct elements $a \in \mathcal{A}$ as suffixes of the $\boldsymbol{\tau}$ substitution words, so there is no non-empty common suffix $\mathbf{s}_1$. Now for any subset $\mathcal{A}^*$ of $\mathcal{A}$, the sequence A^*_n is eventually periodic. If for some $\mathcal{A}^*$, the suffixes of the $\boldsymbol{\tau}$-substitution words of elements in A^*_n are non-empty infinitely often, then the $\boldsymbol{\tau}^n$ words of elements in $\mathcal{A}^*$ will have a prefix like that in Equation 3.1, leading to an element $\boldsymbol{y}$ satisfying $\mathbf{e}^*_M\mathbf{s}^*_1\boldsymbol{\tau}(\boldsymbol{y}) = \boldsymbol{y}$. If $\boldsymbol{y} = \boldsymbol{u}$, then $\mathbf{e}^*_M\mathbf{s}^*_1\boldsymbol{\tau}(\boldsymbol{u}) = \boldsymbol{\tau}(\boldsymbol{u})$ and recognizability implies that $\mathbf{e}^*_M\mathbf{s}^*_1 = \boldsymbol{\tau}(\boldsymbol{u}_{[0,r]})$ for some non-negative r. Using recognizability again, and injectivity, we conclude that $\boldsymbol{u}$ is periodic, a contradiction. This proves necessity. Conversely, letting $\mathcal{A}^* = \mathcal{A}$, and using Proposition 6, $\boldsymbol{u}$ is non-σ-invertible. If $\boldsymbol{y}$ were another branch point, Proposition 7 implies that there are letters a and b, and a sequence $n_j \to \infty$, with $\boldsymbol{y} \in \sigma^{k_j}(\boldsymbol{\tau}^{n_j}(a)) \cap \sigma^{l_j}(\boldsymbol{\tau}^{n_j}(b))$ and $\sigma^{k_j}(\boldsymbol{\tau}^{n_j}(a))$ a maximal common suffix of $\boldsymbol{\tau}^{n_j}(a)$ and $\boldsymbol{\tau}^{n_j}(b)$. Letting $\mathcal{A}^* = \{a, b\}$, we have a contradiction. Thus $\boldsymbol{\tau}$ is quasi-invertible. ■

If $\boldsymbol{\tau}$ is not left proper, then a modified version of Lemma 9 exists. Namely, if there exists an m such that whenever $\mathcal{A}^* \subset \mathcal{A}$ is such that $\{\boldsymbol{\tau}^m(a) : a \in \mathcal{A}^*\}$ has $\boldsymbol{s}^*_1 \neq \emptyset$, then the sequences $\{\mathbf{e}^*_k\boldsymbol{s}^*_1\}$ are all eventually identical, and if for all fixed points $\boldsymbol{u}$, there is a unique a with $a\boldsymbol{u} \in X_{\boldsymbol{\tau}}$, then $\boldsymbol{\tau}$ is quasi-invertible, with branch point $\boldsymbol{y} \neq \boldsymbol{u}$. There is then the possibility that $\boldsymbol{\tau}$ is quasi-invertible, but the set of $\boldsymbol{\tau}$-substitution words do not have a proper maximal common suffix, and this is an obstacle to extending Lemma 10 below to non-left proper substitutions.

If the left proper $\boldsymbol{\tau}$ is such that $\mathbf{e}_n = \mathbf{e}_M$ for $n \geq M$, then the branch point $\mathbf{y}$ satisfies $\mathbf{e}_M\mathbf{s}_1\boldsymbol{\tau}(\boldsymbol{y}) = \boldsymbol{y}$. In the next lemma, we assume that $M = 1$.

LEMMA 10. *Suppose that the reasonable* $\boldsymbol{\tau}$ *is defined as* $\boldsymbol{\tau}(a_i) = \mathbf{p}_i x_i \mathbf{s}_1$, *where* $\{x_1, \dots, x_n\}$ *has at least two distinct elements, and where* $\mathbf{e}_k = \emptyset$ *for* $k \geq 1$. *Define the substitution* $\boldsymbol{\tau}^*$ *as* $\boldsymbol{\tau}^*(a_i) = \mathbf{s}_1\mathbf{p}_i x_i$.

(i) *For each* $n \geq 0$, *and each* $\boldsymbol{w} \in \mathcal{L}_{\boldsymbol{\tau}}$, $\boldsymbol{\tau}^*(\boldsymbol{\tau}^n(\boldsymbol{w}))\mathbf{s}_1 = \mathbf{s}_1\boldsymbol{\tau}^{n+1}(\boldsymbol{w})$.

(ii) *If* $\mathbf{s}_n$ *is as in Lemma 9, then for each* $n \geq 2$, $\mathbf{s}_n = \boldsymbol{\tau}^*(\mathbf{s}_{n-1})\,\mathbf{s}_1$.

(iii) $\mathbf{s}_n \subset (\boldsymbol{\tau}^*)^{n-1}(\mathbf{s}_1)$ *for each* $n \geq 1$.

Proof.

1. We prove this statement only for $\boldsymbol{w} = a_1$; the proof for other one-letter words is the same, and the proof for longer words follows by concatenation. Note that
$$\boldsymbol{\tau}^*(a_1)\,\mathbf{s}_1 = \mathbf{s}_1\mathbf{p}_1 x_1 \mathbf{s}_1 = \mathbf{s}_1\boldsymbol{\tau}(a_1)\,.$$
If the assertion is true for n, then
$$\boldsymbol{\tau}^*(\boldsymbol{\tau}^{n+1}(a_1))\mathbf{s}_1 = \boldsymbol{\tau}^*(\boldsymbol{\tau}^n(\boldsymbol{\tau}(a_1)))\mathbf{s}_1 = \boldsymbol{\tau}^*(\boldsymbol{\tau}^n(\mathbf{p}_1 x_1 \mathbf{s}_1))\mathbf{s}_1 = \mathbf{s}_1\boldsymbol{\tau}^{n+1}(\boldsymbol{\tau}(a_1)) = \mathbf{s}_1\boldsymbol{\tau}^{n+2}(a_1)\,.$$
2. First by Lemma 9 and Part 1 of this lemma, $\mathbf{s}_2 = \mathbf{s}_1\boldsymbol{\tau}(\mathbf{s}_1) = \boldsymbol{\tau}^*(\mathbf{s}_1)\mathbf{s}_1$. Assuming that $\mathbf{s}_n = \boldsymbol{\tau}^*(\mathbf{s}_{n-1})\mathbf{s}^{\mathbf{1}}$, we have
$$\mathbf{s}_{n+1} \overset{Lemma9}{=} \mathbf{s}_n\boldsymbol{\tau}^n(\mathbf{s}_1) \overset{IH}{=} \boldsymbol{\tau}^*(\mathbf{s}_{n-1})\mathbf{s}_1\boldsymbol{\tau}^n(\mathbf{s}_1) \overset{Part1}{=} \boldsymbol{\tau}^*(\mathbf{s}_{n-1})\boldsymbol{\tau}^*(\boldsymbol{\tau}^{n-1}(\mathbf{s}_1))\mathbf{s}_1 = \boldsymbol{\tau}^*(\mathbf{s}_n)\mathbf{s}_1.$$
3. The case $n = 1$ is clear. Assuming $\mathbf{s}_n \subset (\boldsymbol{\tau}^*)^{n-1}(\mathbf{s}_1)$,
$$\mathbf{s}_{n+1} \overset{Part2}{=} \boldsymbol{\tau}^*(\mathbf{s}_n)\mathbf{s}_1 \overset{IH}{\subset} \boldsymbol{\tau}^*((\boldsymbol{\tau}^*)^{n-1})(\mathbf{s}_1))\mathbf{s}_1 = (\boldsymbol{\tau}^*)^n(\mathbf{s}_1)\,\mathbf{s}_1 \subset (\boldsymbol{\tau}^*)^n(\mathbf{s}_1)\,.$$

■

Lemma 10 can be modified if $M > 1$: instead of considering $\boldsymbol{\tau}$, work with $\boldsymbol{\tau}^M$, and as in the lemma, find $\boldsymbol{\tau}^*$ such that $(X_{\boldsymbol{\tau}^*}, \sigma) = (X_{\boldsymbol{\tau}^N}, \sigma)$; now use the fact that $(X_{\boldsymbol{\tau}^N}, \sigma) = (X_{\boldsymbol{\tau}}, \sigma)$.

Corollary 11.
If the left proper, reasonable substitution subshift $(X_{\boldsymbol{\tau}}, \sigma)$ *is quasi-invertible, where* τ *is defined on* $\{a_1, \dots a_n\}$ *with branch point* $\boldsymbol{y}$, *then there exists a left proper, recognizable, minimal, aperiodic, injective, quasi-invertible substitution subshift with* $\boldsymbol{\tau}^*(\boldsymbol{y}) = \boldsymbol{y}$, *and* $(X_{\boldsymbol{\tau}}, \sigma) = (X_{\boldsymbol{\tau}^*}, \sigma)$.

Proof. If $\boldsymbol{\tau}(\boldsymbol{y}) = \boldsymbol{y}$, then let $\boldsymbol{\tau} = \boldsymbol{\tau}^*$. If not, then if necessary considering a power of $\boldsymbol{\tau}$ so that $\mathbf{e}_k = \emptyset$ for $k \geq 1$, let $\boldsymbol{\tau}^*$ be defined as in Lemma 10. The branch point is $\boldsymbol{y} := \lim_{n\to\infty} \mathbf{s}_n$; that it exists and has at least two σ-preimages follows from Lemma 9. That it is $\boldsymbol{\tau}^*$-fixed follows from part 3 of Lemma 10. Thus $X_{\boldsymbol{\tau}^*} \subset X_{\boldsymbol{\tau}}$, and by minimality, this inclusion is an equality. ■

Examples All examples are left proper, and reasonable.

1. If $\boldsymbol{\tau}(a) = aab$, and $\boldsymbol{\tau}(a) = abb$, then $\mathbf{s}_1 = \{b\}$, $\mathbf{e}_k = \emptyset$ for all $k \geq 2$. (In fact if $\boldsymbol{\tau}$ is any reasonable substitution on a two letter alphabet whose substitution words have a proper maximal non-empty suffix, then it is always the case that $\mathbf{e}_k$ is trivial for $k \geq 2$.) The branch point $\boldsymbol{y}$ satisfies $b\boldsymbol{\tau}(\boldsymbol{y}) = \boldsymbol{y}$.
2. The *Tribonnaci substitution* $\boldsymbol{\tau}(1) = 12$, $\boldsymbol{\tau}(2) = 13$, $\boldsymbol{\tau}(3) = 1$ is suffix permutative, so the fixed point $\boldsymbol{u}$ is the only branch point.
3. If $\boldsymbol{\tau}(a) = acb$, $\boldsymbol{\tau}(b) = aba$ and $\boldsymbol{\tau}(c) = aaa$, then only $\boldsymbol{u}$ is a branch point. For, if $\mathcal{A}^* = \{b, c\}$, then the maximal common suffix of $\{\boldsymbol{\tau}^n(b), \boldsymbol{\tau}^n(c)\}$ is $\boldsymbol{\tau}^{n-2}(a)\boldsymbol{\tau}^{n-1}(a)$, and this converges to $\boldsymbol{u}$ as $n \to \infty$.
4. If $\boldsymbol{\tau}(a) = acb$, $\boldsymbol{\tau}(b) = aba$ and $\boldsymbol{\tau}(c) = aca$, then $\{a, b\}$ is a suffix cycle, so $\boldsymbol{u}$ is a branch point. Letting $\mathcal{A}^* = \{b, c\}$, we obtain a branch point $\boldsymbol{y}$ satisfying $a\boldsymbol{\tau}(\boldsymbol{y}) = \boldsymbol{y}$, and $\boldsymbol{y} \neq \boldsymbol{u}$.
5. If $\boldsymbol{\tau}(a) = abc$, $\boldsymbol{\tau}(b) = aacc$, and $\boldsymbol{\tau}(c) = abcc$, then $\mathbf{s}_1 = c$, $A_1 = \{b, c\}$, and for $k \geq 1$, $\mathbf{e}_{2k+1}\mathbf{s}_1 = c$, $A_{2k+1} = \{b, c\}$, $\mathbf{e}_{2k}\mathbf{s}_1 = cc$ and $A_{2k} = \{a, b\}$, so that $(X_{\boldsymbol{\tau}}, \sigma)$ has two branch points, $\boldsymbol{y} = \lim_{n\to\infty} \prod_{k=0}^{n-1} \boldsymbol{\tau}^{2k}(c)\boldsymbol{\tau}^{2k+1}(cc)$ and $\boldsymbol{y}^* =$

$\lim_{n\to\infty}\prod_{k=0}^{n-1}\boldsymbol{\tau}^{2k}(cc)\boldsymbol{\tau}^{2k+1}(c)$; if these were equal then $\boldsymbol{\tau}(c)=c^k$ for some k, a contradiction.

(6) If $\boldsymbol{\tau}(a)=abd$, $\boldsymbol{\tau}(b)=abcd$, $\boldsymbol{\tau}(c)=accd$ and $\boldsymbol{\tau}(d)=aacd$, then for $k\geq 2$, $\mathbf{e}_k\mathbf{s}_1=cd$, $A_k=\{b,c\}$, so that $(X_{\boldsymbol{\tau}},\sigma)$ has exactly one branch point, satisfying $cd\boldsymbol{\tau}(\boldsymbol{y})=\boldsymbol{y}$.

(7) If $\boldsymbol{\tau}(a)=abad$, $\boldsymbol{\tau}(b)=ab$, $\boldsymbol{\tau}(c)=ad$, and $\boldsymbol{\tau}(d)=abc$ is recognizable: although $\boldsymbol{\tau}(c)$ is a proper suffix of $\boldsymbol{\tau}(a)$, a is always followed by b or d in X, while c is only followed by a. So $\boldsymbol{\tau}(a)$ is always followed by $\boldsymbol{\tau}(b)$ or $\boldsymbol{\tau}(d)$, and $\boldsymbol{\tau}(c)$ is always followed by $\boldsymbol{\tau}(a)$. Hence long words satisfying Part 2 of Theorem 3 do not exist.
As $\{b,c,d\}$ belong to suffix cycles, $\boldsymbol{u}$ has three σ-preimages. If $\mathcal{A}^*=\{a,c\}$ then the words $\{\boldsymbol{\tau}^n(a),\boldsymbol{\tau}^n(c)\}$ have $\boldsymbol{\tau}^n(c)$ as a maximal common suffix, and these words also converge to $\boldsymbol{u}$. Hence $\boldsymbol{\tau}$ is quasi-invertible.

(8) If $\boldsymbol{\tau}(a)=aad$, $\boldsymbol{\tau}(b)=abd$, $\boldsymbol{\tau}(c)=ac$, and $\boldsymbol{\tau}(d)=abc$. All $\boldsymbol{\tau}^2$-substitution words have c as a suffix, so $\boldsymbol{u}$ is σ-invertible. Also, $\mathbf{e}_k=dab$ for $k\geq 2$, and $A_k=\{a,b\}$ for all k. Working with $\boldsymbol{\tau}^4$, there is only one branch point $\boldsymbol{y}$; it satisfies $dabc\,\boldsymbol{\tau}(\boldsymbol{y})=\boldsymbol{y}$.

Unilateral recognizability is less amenable than bilateral recognizability, which all aperiodic primitive substitutions possess, and for whom the sequence of partitions

$$\mathcal{P}_n^{'}:=\{[\sigma^k(\boldsymbol{\tau}^n(a))]:a\in\mathcal{A},\ \ 0\leq k<|\tau^n(a)|\}$$

is a nested sequence which spans the topology of $(X_{\mathbb{Z}}(\boldsymbol{\tau}),\sigma)$ (see Proposition 14, [**DHS99**]). In the one sided subshift $(X_{\boldsymbol{\tau}},\sigma)$, the sets in $\mathcal{P}_n^{'}$ need not be disjoint. For example, if $\boldsymbol{\tau}$ has a branch point $\boldsymbol{y}\neq\boldsymbol{u}$, then, for each n, there are letters a, b, and numbers n_1 and n_2 such that $|\boldsymbol{\tau}^n(a)|-n_1=|\boldsymbol{\tau}^n(b)|-n_2$ and $\sigma^{n_1}(\boldsymbol{\tau}^n(a))=\sigma^{n_2}(\boldsymbol{\tau}^n(b))$, while $\sigma^{n_1-1}(\boldsymbol{\tau}^n(a))\neq\sigma^{n_2-1}(\boldsymbol{\tau}^n(b))$. If $\boldsymbol{\tau}$ is such that $\boldsymbol{u}$ is the only element in $X_{\boldsymbol{\tau}}$ with several σ-preimages, and we modify the definition of $\mathcal{P}_n^{'}$ slightly this does not happen. Define

$$\mathcal{P}_n:=\{[\sigma^k(\boldsymbol{\tau}^n(a))]\cap\sigma^{-|\sigma^k(\boldsymbol{\tau}^n(a))|}\boldsymbol{\tau}(X):a\in\mathcal{A},\ \ 0\leq k<|\tau^n(a)|\}\,.$$

If $\boldsymbol{w}=\sigma^k(\boldsymbol{\tau}^n(a))$, let $[\boldsymbol{w}]_*=[\sigma^k(\boldsymbol{\tau}^n(a))]\cap\sigma^{-|\sigma^k(\boldsymbol{\tau}^n(a))|}\boldsymbol{\tau}(X)$.

COROLLARY 12. *Let $\boldsymbol{\tau}$ be left proper and reasonable, and suppose that $\boldsymbol{u}$ is the only branch point in $X_{\boldsymbol{\tau}}$.*
Then

(i) *For every $n>0$, $\mathcal{P}_n$ is a clopen partition, and the sequence of bases is decreasing.*
(ii) *For each n, $\mathcal{P}_{n+1}$ is a refinement of $\mathcal{P}_n$: i.e. every element of $\mathcal{P}_{n+1}$ is contained in an element of $\mathcal{P}_n$.*
(iii) *The intersection of the bases of $\mathcal{P}_n$ consists of a unique point, and the sequence $\{\mathcal{P}_n\}$ spans the topology of (X,σ).*

Proof.

(1) Part 2 of Proposition 4 and the compactness of $X_{\boldsymbol{\tau}}$ imply that the elements in $\mathcal{P}_n$ are clopen. To show that they are disjoint, suppose that $\boldsymbol{y}\in[\boldsymbol{w}]_*\cap[\boldsymbol{w}']_*$ where $\boldsymbol{w}=\sigma^k(\tau^n(a))$ and $\boldsymbol{w}'=\sigma^l(\boldsymbol{\tau}^n(b))$. If $|\boldsymbol{w}|<|\boldsymbol{w}'|$ (or $|\boldsymbol{w}|>|\boldsymbol{w}'|$), then $\sigma^{|\boldsymbol{w}|}(\boldsymbol{y})$ (or $\sigma^{|\boldsymbol{w}'|}(\boldsymbol{y})$) has two representations of the form $\sigma^j(\boldsymbol{\tau}(\boldsymbol{x}))$, a contradiction, using Corollary 5. So $|\boldsymbol{w}|=|\boldsymbol{w}'|$, which means that $\boldsymbol{w}$ is a common suffix of $\boldsymbol{\tau}^n(a)$ and $\boldsymbol{\tau}^n(b)$. Since $\boldsymbol{\tau}^n(a)\neq\boldsymbol{\tau}^n(b)$, these two words have a maximal proper common suffix, say $\sigma^{k'}(\tau^n(a))=\sigma^{l'}(\boldsymbol{\tau}^n(b))$, and here one of k' or l' must be positive - say k'. Thus $\sigma^{-(k-k')}(\boldsymbol{y})=\sigma^{-(l-l')}(\boldsymbol{y})$ has two preimages, so by assumption $\boldsymbol{u}=\sigma^{-(k-k')}\boldsymbol{y}\in\sigma^{-(k-k')}[\sigma^k(\boldsymbol{\tau}^n(a))]_*=[\sigma^{k'}\boldsymbol{\tau}^n(a)]_*$, once again a contradiction to Corollary 5.
(2) The proof of Parts 2 and 3 are similar to the proof of Proposition 14, [**DHS99**]. We repeat the argument here. Take an element $[\boldsymbol{w}]_*\in\mathcal{P}_{n+1}$: we need to show that $[\boldsymbol{w}]_*\subset[\boldsymbol{w}']_*$ where $[\boldsymbol{w}']^*\in\mathcal{P}_n$. Suppose that $\boldsymbol{w}=\sigma^k(\boldsymbol{\tau}^{n+1}(a))$, where $0<k<$

$|\boldsymbol{\tau}^{n+1}(a)|$. Write $\boldsymbol{\tau}^{n+1}(a) = \boldsymbol{\tau}^n(a_1)\,\boldsymbol{\tau}^n(a_2)\dots\boldsymbol{\tau}^n(a_m)$ where $\boldsymbol{\tau}(a) = a_1\,a_2\dots a_m$. For some j, $|\boldsymbol{\tau}^n(a_1\dots a_j)| \leq k < |\boldsymbol{\tau}^n(a_1\dots a_{j+1})|$, so let $l := k - |\boldsymbol{\tau}^n(a_1\dots a_j)|$. Then $\sigma^k(\boldsymbol{\tau}^{n+1}([a])) \subset \sigma^l\boldsymbol{\tau}^n[a']$, where $a' = a_{j+1}$.

(3) Given m positive, we show that for all n large, each element $\mathbf{p}$ of $\mathcal{P}_n$ is contained in some $\mathbf{c}$ where $\mathbf{c}$ is a word of length $m+1$. Suppose that all $\boldsymbol{\tau}$-substitution words start with l. Write $l_n = |\boldsymbol{\tau}^{n-1}(l)|$. Choose n so that $l_n > m$ (Minimality implies that $\lim_{n\to\infty} l_n = \infty$).

Fix $\boldsymbol{w} \in \mathcal{W}_n$, and suppose that $\boldsymbol{w} = \sigma^k(\boldsymbol{\tau}^n(a))$ for some $a \in \mathcal{A}$ and $k \in [0, |\boldsymbol{\tau}^n(a)|)$. If $\boldsymbol{y} \in [\boldsymbol{w}]^*$ then there is some $\boldsymbol{x} \in [a]$ with $\boldsymbol{y} = \sigma^k\boldsymbol{\tau}^n(\boldsymbol{x})$. Since $\boldsymbol{\tau}$ is left proper, $\boldsymbol{\tau}^n(a)\boldsymbol{\tau}^{n-1}(l)$ is a prefix of $\boldsymbol{\tau}^n(\boldsymbol{x})$, and so $\boldsymbol{y} \in [\sigma^k\boldsymbol{\tau}^n(a)\boldsymbol{\tau}^{n-1}(l)]$, and this last cylinder set has length greater than m. So, $\boldsymbol{y}$ starts with a block of length m depending only on $\boldsymbol{w}$ and not on $\boldsymbol{y}$. ■

4. *Bratteli Diagrams*

A *Bratteli diagram* $\mathcal{B} = (\mathcal{V}, \mathcal{E})$ is an infinite directed graph with *vertex set* $\mathcal{V} = \bigsqcup_{n=0}^{\infty}\mathcal{V}_n$ and *edge set* $\mathcal{E} = \bigsqcup_{n=1}^{\infty}\mathcal{E}_n$, where all $\mathcal{V}_n$'s and $\mathcal{E}_n$'s are finite, $\mathcal{V}_0 = \{v_0\}$, and, if x is an edge in $\mathcal{E}_n$, the *source* $s(x)$ of x lies in $\mathcal{V}_{n-1}$ and the *range* $r(x)$ of x lies in $\mathcal{V}_n$. We assume that $s^{-1}(v) \neq \emptyset$ for each $v \in \mathcal{V}$ and $r^{-1}(v) \neq \emptyset$ for all $v \in \mathcal{V}\backslash v_0$. We will use $x, y\dots$ when referring to edges, and $a, b, \dots$ when referring to vertices.

A finite set of edges $\{x_{n+k}\}_{k=1}^{K}$, with $s(x_{n+k+1}) = r(x_{n+k})$ for $1 \leq k \leq K-1$, is called a *path* from $s(x_{n+1})$ to $r(x_{n+K})$. Similarly an *infinite path* in $\mathcal{B}$ is a sequence $\boldsymbol{x} = \{x_n\}_{n=\infty}^{1}$, with $x_n \in \mathcal{E}_n$ for $n \geq 1$, and $s(x_{n+1}) = r(x_n)$ for $n \geq 1$. We write $\dots a_{n+1} \overset{x_{n+1}}{\leftarrow} a_n \overset{x_n}{\leftarrow} \dots \overset{x_2}{\leftarrow} a_1 \overset{x_1}{\leftarrow} v_0$ where $r(x_n) = a_n$, or $(\mathbf{a}, \boldsymbol{x}) = \{(a_n, x_n)\}_{n\geq 1}$, when referring to an element $\boldsymbol{x}$ in $X_{\mathcal{B}}$. The set of all infinite paths in $\mathcal{B}$ will be denoted $X_{\mathcal{B}}$ (a subset of $\Pi_{n\geq1}\mathcal{E}_n$), and $X_{\mathcal{B}}$ is endowed with the topology induced from the product topology on $\Pi_{n\geq1}\mathcal{E}_n$. Thus $X_{\mathcal{B}}$ is a compact metric space.

Two Bratteli diagrams $\mathcal{B} = (\mathcal{V}, \mathcal{E})$ and $\mathcal{B}' = (\mathcal{V}', \mathcal{E}')$ are *isomorphic* if there exists a pair of bijections $f_{\mathcal{V}} : \mathcal{V} \to \mathcal{V}'$ and $f_{\mathcal{E}} : \mathcal{E} \to \mathcal{E}'$ satisfying $f_{\mathcal{V}}(a) \in \mathcal{V}'_n$ if $a \in \mathcal{V}_n$, and $s(f_{\mathcal{E}}(x)) = f_{\mathcal{V}}(s(x))$, $r(f_{\mathcal{E}}(x)) = f_{\mathcal{V}}(r(x))$ whenever $x \in \mathcal{E}$. Let $\{n_k\}_{k=0}^{\infty}$ be a sequence of increasing integers with $n_0 = 0$. Then $\mathcal{B}' = (\mathcal{V}', \mathcal{E}')$ is a *telescoping* of $\mathcal{B} = (\mathcal{V}, \mathcal{E})$ if $\mathcal{V}'_k = \mathcal{V}_{n_k}$ (with the vertex $v \in \mathcal{V}_{n_k}$ labelled as $v' \in \mathcal{V}'_k$), and the number of edges from $v'_k \in \mathcal{V}'_k$ to $v'_{k+1} \in \mathcal{V}'_{k+1}$ is the number of paths from $v_k \in \mathcal{V}_{n_k}$ to $v_{k+1} \in \mathcal{V}_{n_{k+1}}$. Conversely, we can perform a *splitting* by introducing a new level between two consecutive levels $\mathcal{V}_{n-1}$ and $\mathcal{V}_n$, so that the number of new vertices equals the edges in $\mathcal{E}_n$, and each vertex in the new level is the source and range of exactly one edge. We consider two Bratteli diagrams $\mathcal{B}$ and $\mathcal{B}'$ *equivalent* if $\mathcal{B}'$ can be obtained from $\mathcal{B}$ by isomorphism, telescoping and splitting. Thus when we talk about a Bratteli diagram we are talking about an equivalence class of diagrams.

We say that $\mathcal{B}$ is *simple* if there exists a telescoping $\mathcal{B}' = (\mathcal{V}', \mathcal{E}')$ of $\mathcal{B}$ so that for any $a \in \mathcal{V}'_n$ and $b \in \mathcal{V}'_{n+1}$, there is at least one edge from a to b. If $\boldsymbol{x} = \{x_n\}_{n=\infty}^{1}$ and $\boldsymbol{x}' = \{x'_n\}_{n=\infty}^{1}$ are two elements in $X_{\mathcal{B}}$, we write $\boldsymbol{x} \sim \boldsymbol{x}'$ if the tails of $\boldsymbol{x}$ and $\boldsymbol{x}'$ are equal. It follows that $\sim$ is an equivalence relation, and if $\mathcal{B}$ is simple, then each equivalence class for $\sim$ is dense in $X_{\mathcal{B}}$, and $X_{\mathcal{B}}$ has no isolated points, making it a Cantor space.

4.0.1. *Ordering $X_{\mathcal{B}}$* Let $n \geq 1$. For each $a \in \mathcal{V}_n$, let $\mathcal{E}_n(a) = \{x \in \mathcal{E}_n : r(x) = a\}$. We say $\mathcal{B}$ is *ordered* if there is a linear order $\geq$ on each $\mathcal{E}_n(a)$; elements of $\mathcal{E}_n(a)$ will then be labelled $1, 2, \dots$ according to their order. If $a \in \mathcal{V}\backslash\{v_0\}$, define $|a| := |\mathcal{E}_n(a)|$, so that $\mathcal{E}_n(a) = \{1, 2, \dots |a|\}$.

The linear order on edges in each $\mathcal{E}_n(a)$ induces a partial ordering on paths from $\mathcal{V}_m$ to $\mathcal{V}_n$: the two paths $\boldsymbol{x} = a_n \overset{x_n}{\leftarrow} a_{n-1} \overset{x_{n-1}}{\leftarrow} \dots \overset{x_{m+1}}{\leftarrow} a_m$ and $\mathbf{x}' = a'_n \overset{x'_n}{\leftarrow} a'_{n-1} \overset{x'_{n-1}}{\leftarrow} \dots \overset{x'_{m+1}}{\leftarrow} a'_m$ from $\mathcal{E}_m$ to $\mathcal{E}_n$ are comparable with $\boldsymbol{x} < \boldsymbol{x}'$ if there is some $k \in [m+1, n]$ with $x_k < x'_k$ and $x_j = x'_j$ for $k+1 \leq j \leq n$.

Finally, two elements $\boldsymbol{x}$, $\boldsymbol{x}' \in X_{\mathcal{B}}$ are *comparable* with $\boldsymbol{x} < \boldsymbol{x}'$ if there is a k such that $x_n = x'_n$ for all $n > k$, and $x_k < x'_k$. Thus each equivalence class for $\sim$ is ordered. There is the obvious notion of ordered isomorphism of two ordered Bratteli diagrams $\mathcal{B}$, $\mathcal{B}'$: the isomorphism between $\mathcal{B}$ and $\mathcal{B}'$ also has to satisfy $f_{\mathcal{E}}(x) \leq f_{\mathcal{E}}(y)$ if $x \leq y$. If $\mathcal{B}'$ is a telescoping of the ordered Bratteli diagram $\mathcal{B}$, then the order induced on $\mathcal{B}'$ from the order on $\mathcal{B}$ makes $\mathcal{B}'$ an ordered Bratteli diagram. We say that the ordered Bratteli diagrams $\mathcal{B}$, $\mathcal{B}'$ are *equivalent* if $\mathcal{B}'$ is the image of $\mathcal{B}$ by telescoping and order isomorphism.

An infinite path is *maximal* (*minimal*) if all the edges making up the path are maximal (minimal). If $\boldsymbol{x} = \{x_n\}_{n=\infty}^{1}$ is not maximal, let k be the smallest integer such that $x_k \in \mathcal{E}_k(a_k)$ is not a maximal edge, and let y_k be the successor of x_k in $\mathcal{E}_k(a_k)$. Then the *successor* $\mathcal{V}_{\mathcal{B}}(\boldsymbol{x})$ is defined to be $\mathcal{V}_{\mathcal{B}}(\boldsymbol{x}) = \dots x_{k+2}\, x_{k+1}\, y_k\, 0 \dots 0$. Similarly, every non-minimal path has a unique *predecessor*. Let $X_{\min}$, $(X_{\max}) \subset X_{\mathcal{B}}$ be defined as the set of minimal (maximal) elements of $X_{\mathcal{B}}$. By compactness, these sets are non empty. Simple ordered Bratteli diagrams which have a unique minimal and maximal element (called $\boldsymbol{x}_{\min}$ and $\boldsymbol{x}_{\max}$ respectively) are called *proper*, and those with a unique minimal element are called *semi-proper*. If $\mathcal{B}$ is semi-proper, then $V_{\mathcal{B}}$ can be extended to a continuous surjection on $X_{\mathcal{B}}$ by setting $V_{\mathcal{B}}(\boldsymbol{x}_{\max}) = \boldsymbol{x}_{\min}$, which is a homeomorphism if $\mathcal{B}$ is proper. We call $(X_{\mathcal{B}}, V_{\mathcal{B}})$ a *Bratteli-Vershik* or *adic* system. Note that $(X_{\mathcal{B}}, V_{\mathcal{B}})$ is a minimal system, since $V_{\mathcal{B}}$ orbits are equivalence classes for $\sim$. Let us say that two Cantor systems $(X_i, T_i, \boldsymbol{x}_i)$, $i = 1, 2$ are *pointedly isomorphic* if there exists a homeomorphism $f : X_1 \to X_2$ with $f \circ T_1 = T_2 \circ f$ and $f(x_1) = x_2$. The adic system associated to an equivalence class of ordered semi-proper Bratteli diagrams is well defined up to pointed isomorphism. This was proved in Section 4, [**HPS92**] for proper Bratteli diagrams; the proof is similar for semi-proper Bratteli diagrams:

Proposition 13. *Let $\mathcal{B}$ and $\mathcal{B}'$ be semi-proper ordered Bratteli diagrams, with the same number of maximal elements. Then $\mathcal{B}$ is equivalent to $\mathcal{B}'$ if and only if $(X_{\mathcal{B}}, V_{\mathcal{B}}, \boldsymbol{x}_{\min})$ is pointedly isomorphic to $(X_{\mathcal{B}'}, V_{\mathcal{B}'}, \boldsymbol{x}'_{\min})$.* ■

Let $\boldsymbol{\tau}$ be a primitive substitution on $\mathcal{A}$. The *Bratteli diagram associated with $\boldsymbol{\tau}$* has vertex sets $\mathcal{V}_n = \mathcal{A}$, for each $n \geq 1$. There is exactly one edge from each vertex in $\mathcal{V}_1$ to v_0. If $\boldsymbol{\tau}(a) = a_1 a_2 \dots a_n$, then there is an edge from vertex b in $\mathcal{V}_{n-1}$ to vertex $a \in \mathcal{V}_n$, and it is labelled i, if and only if $a_i = b$. If $1 \leq m < n$, then the number of paths in $\mathcal{B}$ from (a, n) to (b, m) is the number of occurrences of b in $\boldsymbol{\tau}^{(n-m)}(a)$. Primitivity implies that there is a positive k such that for any two letters a and b, there is at least one path from $(a, n+k)$ to (b, n). The Bratteli diagram $\mathcal{B}' = (\mathcal{V}', \mathcal{E}')$ for $\boldsymbol{\tau}^k$ is the telescoping of the Bratteli diagram $\mathcal{B} = (\mathcal{V}, \mathcal{E})$ for $\boldsymbol{\tau}$, with $\mathcal{V}'_n = \mathcal{V}_{nk}$ for $n > 1$ and $\mathcal{V}'_1 = \mathcal{V}_1$. An example of a Bratteli diagram is illustrated in Figure 1.

The connection between adic systems and one sided left proper reasonable substitutions is given by the next result. It is the appropriate generalization of [**DHS99**, Prop 20] to one sided substitution systems.

Theorem 14.



FIGURE 1. *The Bratteli diagram associated with the substitution* $\boldsymbol{\tau}(a) = abb, \ \ \boldsymbol{\tau}(b) = ab$.

If $\boldsymbol{\tau}$ *is left proper, reasonable and quasi-invertible, with fixed point* $\boldsymbol{u}$*, there exists a semi-proper, stationary Bratteli diagram* $\mathcal{B}$ *which is semi-proper, and stationary, such that* $(X_{\mathcal{B}}, V_{\mathcal{B}})$ *is topologically conjugate to* (X, σ).

Proof. If $\boldsymbol{u}$ is the branch point, let $\mathcal{B}$ be the Bratteli diagram associated with $\boldsymbol{\tau}$. Since $\boldsymbol{\tau}$ is minimal and left proper, it is primitive. In this case the sequence of sets $\mathcal{P}_n$ defined in Section 3 are a refining sequence of partitions, by Corollary 12, which generate the topology of $(X_{\boldsymbol{\tau}}, \sigma)$. As in the proof of Proposition 16, [**DHS99**], we construct an isomorphism $F : X_{\boldsymbol{\tau}} \to X_{\mathcal{B}}$ by having the path $F(\boldsymbol{y})$ passing through the vertex in $\mathcal{V}_n$ that corresponds to the tower in $\mathcal{P}_n$ where $\boldsymbol{y}$ is located. The proof that F is a conjugacy between $(X_{\mathcal{B}}, V_{\mathcal{B}})$ and $(X_{\boldsymbol{\tau}}, \sigma)$ is as in the aforementioned proof.

If $\boldsymbol{\tau}$ is left proper and the branch point $\boldsymbol{y} \neq \boldsymbol{u}$, work with the substitution $\boldsymbol{\tau}^*$ in Corollary 11. As above, $(X_{\boldsymbol{\tau}^*}, \sigma)$ is conjugate to $(X_{\mathcal{B}^*}, V_{\mathcal{B}^*})$, and since $(X_{\boldsymbol{\tau}}, \sigma) = (X_{\boldsymbol{\tau}^*}, \sigma)$, the result follows. ■

Lemma 15 in [**For97**] is used in the two sided version of this previous result, to show that (aperiodic, primitive) substitution subshifts are conjugate to a stationary adic system where there are only single edges from the vertex v_0 to any vertex in $\mathcal{V}_1$. Although the statement of this lemma is also true for semi-proper Bratelli diagrams, it is not clear that the resulting stationary diagram corresponds to a recognizable, or injective substitution. See for example, Figure 4 in [**DHS99**].

5. *Induced substitutions*

We now introduce concepts needed to extend Theorem 14 to non-left proper substitutions.

5.1. *Induced transformations*

Suppose that $T : X \to X$ is a minimal continuous transformation with X a Cantor space - henceforth called a Cantor system. If $U \subset X$ is a clopen set, the *system* (U, T_U) *induced by* (X, T) *on* U, is defined by $T_U(\boldsymbol{x}) = T^n(\boldsymbol{x})$, where n is the least positive natural number such that $T^n(x) \in U$. T_U is well defined since T is minimal, and since X is compact, n can only take a finite number of values. The Cantor systems (X_1, T_1) and (X_2, T_2) are (topologically) *Kakutani equivalent* ([**PS73**] is first reference of this) if there exist clopen sets $U_i \subset X_i$ such that the respective induced systems are isomorphic. Topological Kakutani equivalence is much more stringent than measurable Kakutani equivalence - for example, all rank one transformations are measurably Kakutani equivalent, but it can be shown, for example, that the *Chacon* substitution $\boldsymbol{\tau}(0) = 0010$, $\boldsymbol{\tau}(1) = 1$ is Kakutani equivalent only to the substitutions $\boldsymbol{\tau}(0) = 001^k0$, $\boldsymbol{\tau}(1) = 1$, or $\boldsymbol{\tau}(0) = 01^k00$, $\boldsymbol{\tau}(1) = 1$.

Conversely, given a dynamical system (U, S), and a continuous function $h : U \to \{1, \dots H\}$ (called a *height function*), we build a new dynamical system (X, T) in the following way. For each k in $\{1, \dots H\}$, let $U_k^0 = h^{-1}(\{k\})$. If $k \in \{2, \dots H\}$ and $1 \leq j \leq k$, let U_k^j be a homeomorphic copy of U_k^0. Let $\phi_k^i : U_k^i \to U_k^{i+1}$ be a homeomorphism for each $k \in \{2, \dots H\}$ and each $i \in \{0, \dots k-2\}$, $\phi_1^0 : U_1^0 \to U_1^0$ be the identity map, and let $X = \cup_{k=1}^{H} \cup_{j=0}^{k-1} U_k^j$. Define $T : X \to X$ as $T(x) = \phi_k^i(x)$ if $x \in U_k^i$ for $i < k-1$, and $T(x) = S((\phi_k^{k-2} \circ \phi_k^{k-3} \circ \dots \circ \phi_k^0)^{-1}(x))$ if $x \in U_k^{k-1}$. If this is the case, we say that (X, T) is a *primitive* of (U, S) (with respect to h). If T is a homeomorphism, and U is any clopen set in X, then (X, T) is a primitive of (U, T_U), where $h(x)$ is the first return time under T of $x \in U$ (see [**PS73**]), but this is not always the case with non-invertible systems. If (X, T) is quasi-invertible, with branch point $y \in U$, then it is possible to recover (X, T) as a primitive of (U, T_U) with respect to the return time function h. Even if $\boldsymbol{y} \notin U$, we will see that (X, T) can sometimes still be recovered, but as a quotient space of the primitive constructed above. In this case, if (U, T_U) has an appropriate adic representation, it is possible to extend this representation to (X, T), as in the proof of Theorem 3.8, in [**GPS95**]:

THEOREM 15. *Let $(X_{\mathcal{B}}, V_{\mathcal{B}})$ be an adic system, where $\mathcal{B}$ is semi proper, simple, and with M maximal elements. Suppose that the aperiodic M-quasi-invertible Cantor system (Y, T), with branch point $\boldsymbol{y}$ has an induced system (Y', S) where $\boldsymbol{y} \in Y'$. If (Y', S) is isomorphic to $(X_{\mathcal{B}}, V_{\mathcal{B}})$, then (Y, T) is isomorphic to $(X_{\mathcal{B}'}, V_{\mathcal{B}'})$ where $\mathcal{B}'$ is obtained from $\mathcal{B}$ by adding or removing a finite number of edges to $\mathcal{E}_1$, and changing the ordering on the affected vertices and edges.* ■

5.2. *Return words*

Let $\boldsymbol{\tau}$ be a (not necessarily left proper) reasonable substitution with fixed point $\boldsymbol{u}$, and suppose that $u_0 = a$. A *return word to a* in $\boldsymbol{u}$ is a word $\boldsymbol{w}$ such that

(1) u_0 is a prefix of $\boldsymbol{w}$;
(2) There is no other occurrence of u_0 in $\boldsymbol{w}$;
(3) $\boldsymbol{w}u_0 \in \mathcal{L}(X)$.

The set of return words $\mathcal{R}$ is finite, since $\boldsymbol{u}$ is almost periodic. This notion is the one-sided generalization of two sided return words in [**DHS99**], and we use their notation here. Ordering $\mathcal{R}$ according to the order of appearance of a return word in $\boldsymbol{u}$, we have a bijection ψ from $R := \{1, 2, \dots |\mathcal{R}|\}$ to $\mathcal{R}$. The $\boldsymbol{\tau}$-fixed point $\boldsymbol{u}$, and so every element in X, is a concatenation of return words, with possibly a suffix of a return word as a prefix. Extend ψ by concatenation to $\psi : R^{\mathbb{N}} \to [u_0] \subset \mathcal{A}^{\mathbb{N}}$. Since any $\boldsymbol{y} \in \psi(R^{\mathbb{N}})$ can only be partitioned in one way, using Property 2 of the definition of a return word, ψ is injective.

If $U = [u_0]$, then $\sigma_U(\boldsymbol{x}) = \sigma^{h(\boldsymbol{x})}(\boldsymbol{x})$ where $h(x)$ is the length of the first return word we see in $\boldsymbol{x}$. In this case, given h and (U, σ_U), the subsets U_k^j of the primitive $X' = \cup_{k=1}^{H} \cup_{j=0}^{k-1} U_k^j$ can be identified with $\sigma^j(U_k^0)$, and the maps ϕ_k^j defined in Section 5.1 can be identified with the shift map σ. If $\boldsymbol{x} \in X_{\boldsymbol{\tau}}$ has M-preimages, and $\boldsymbol{x} \notin U$, then it will have M representations in X' -namely if L is the least positive integer such that $\sigma^L(\boldsymbol{x}) \in U$, and $\boldsymbol{x}_1, \dots \boldsymbol{x}_M$ are the σ-preimages of $\boldsymbol{x}$ then each $\boldsymbol{x}_i \in U_{k_i}^{j_i}$ for distinct $k_i' s$ where $j_i = k_i - L - 1$. So here X' is not an isomorphic copy of $X_{\boldsymbol{\tau}}$. Let $\boldsymbol{x}^1, \dots \boldsymbol{x}^M$ be the M representations of $\boldsymbol{x}$ in X'. Define the equivalence relation $\sim$ on X' where $\boldsymbol{x}^i \sim \boldsymbol{x}^j$ and $\phi_{k_i}^{j_i+l}\boldsymbol{x}^i \sim \phi_{k_j}^{j_i+l}\boldsymbol{x}^j$ for all $i, j \in \{1, \dots M\}$, and $l \in \{1, \dots L-1\}$. All other elements are equivalent only to themselves. Let $X_{\sim}^{'}$ be the collection of equivalence classes of $\sim$, and define $\pi : X' \to X_{\sim}^{'}$ to be the mapping sending $\boldsymbol{x} \in X'$ to its equivalence class. Endow $X_{\sim}^{'}$ with the quotient topology, and let $\sigma_{\sim} : X_{\sim}^{'} \to X_{\sim}^{'}$ be defined in the obvious way.

Lemma 16. *$(X^{'}_\sim, \sigma_\sim)$ is topologically conjugate to $(X_{\boldsymbol{\tau}}, \sigma)$.*

Proof. Let $\phi : X^{'}_\sim \to X_{\boldsymbol{\tau}}$ be defined as $\phi([\boldsymbol{x}^i]_\sim) = \boldsymbol{x}$, $\phi([\phi^{j_i+l}_{k_i}\boldsymbol{x}^i]_\sim) = \sigma^l(\boldsymbol{x})$ when $1 \leq l \leq L-1$ and $\phi(\boldsymbol{y}) = \boldsymbol{y}$ otherwise. It is straightforward to show that $\sigma \circ \phi = \phi \circ \sigma_\sim$. We have to show that ϕ and ϕ^{-1} are continuous. Let U be open in $X^{'}_\sim$. If none of the elements $[\phi^{j_i+l}_{k_i}\boldsymbol{x}^i]_\sim$ are elements of U, then $\phi(U) = U$ and this is open in $X_{\boldsymbol{\tau}}$. If for some l, $[\phi^{j_i+l}_{k_i}\boldsymbol{x}^i]_\sim \in U$, then, since $X_\sim$ has the quotient topology, there are open cylinder sets $V_i \subset X'$ with $\phi^{j_i+l}_{k_i}\boldsymbol{x}^i \in V_i$ and $\pi(\cup_i V_i) \subset U$. Elements in $\pi(\cup_i V_i)$ agree with $\boldsymbol{x}$ on some large initial segment, so $\phi\pi(\cup_i V_i)$ contains a cylinder set containing $\boldsymbol{x}$. Thus $\phi(U)$ is open. Conversely, suppose that $U \subset X_{\boldsymbol{\tau}}$ is open and contains $\sigma^l(\boldsymbol{x})$ for some $l \in \{0, \ldots L-1\}$. Then U contains all points that agree with $\sigma^l(\boldsymbol{x})$ on some large initial segment, so that $\cup_i V_i \subset \pi^{-1}(\phi^{-1}(U))$ where V_i is a cylinder set in $U^{j_i+l}_{k_i}$, and $\sigma^l\boldsymbol{x}^i \in V_i$. Thus $\pi^{-1} \circ \phi^{-1}(U)$ is open. Since $X^{'}_\sim$ has the quotient topology, this means that $\phi^{-1}(U)$ is open. ■

Let $\mathcal{D}(\boldsymbol{u}) \in R^{\mathbb{N}}$ be the unique sequence such that $\psi(\mathcal{D}(\boldsymbol{u})) = \boldsymbol{u}$. Let (Y, σ) be the subshift spanned by $\mathcal{D}(\boldsymbol{u}) \in R^{\mathbb{N}}$.

Lemma 17. *(Y, σ) is isomorphic to the system induced by (X, σ) on $[u_0]$ via the map ψ.*

Proof. For each $\boldsymbol{y} \in Y$, $\psi(\sigma\boldsymbol{y}) = \sigma^k(\psi(\boldsymbol{y}))$, where k is the length of the word in $\mathcal{R}$ corresponding to y_0. So $\psi(\sigma^n(\boldsymbol{y})) \in [u_0]$ for all natural n, thus $\psi(Y) \subset [u_o]$. If $\boldsymbol{x} \in [u_0]$, then $\sigma^{n_k}(\boldsymbol{u}) \to \boldsymbol{x}$ for some $n_k \to \infty$. For large k, $\sigma^{n_k}(\boldsymbol{u}) \in [u_0]$, so $\sigma^{n_k}(\boldsymbol{u}) = \psi(\sigma^{j_k}(\mathcal{D}(\boldsymbol{u})) \in \psi(Y)$ for some $j_k \to \infty$. Hence $\boldsymbol{x} \in \psi(Y)$. It follows that $\psi \circ \sigma = \sigma_{[u_0]} \circ \psi$ for $y \in Y$. ■

The substitution $\boldsymbol{\tau}$ can be used to define a substitution on R: if $\boldsymbol{w}$ is a return word, then $\boldsymbol{w}$ starts with u_0, and since $\boldsymbol{\tau}(u_0)$ starts with u_0, so does $\boldsymbol{\tau}(\boldsymbol{w})$. Also, $\boldsymbol{w}u_0$ is a word in $\boldsymbol{u}$, so $\boldsymbol{\tau}(\mathbf{w}u_0) = \boldsymbol{\tau}(\boldsymbol{w})u_0\ldots$ is also a word in $\boldsymbol{u}$. Hence $\boldsymbol{\tau}(\boldsymbol{w})u_0\ldots$ is a concatenation of unique return words, and a prefix of some return word, so that $\boldsymbol{\tau}(\boldsymbol{w})$ is a unique concatenation of return words. Thus if $j \in R$ corresponds to $\boldsymbol{w} \in \mathcal{R}$, and $\boldsymbol{\tau}(\boldsymbol{w}) = \boldsymbol{w}_1\boldsymbol{w}_2\ldots\boldsymbol{w}_r$, define $\boldsymbol{\tau}_1(j) := i_1 i_2 \ldots i_r$ where $i_k \in R$ corresponds to $\boldsymbol{w}_k \in \mathcal{R}$.

Lemma 18. *Suppose $\boldsymbol{\tau}$ is reasonable with generating fixed point $\boldsymbol{u}$. Then*

(i) *$\boldsymbol{\tau}_1$ (or some power of $\boldsymbol{\tau}_1$) is left proper, minimal, injective, with $\mathcal{D}(\boldsymbol{u})$ its generating fixed point;*
(ii) *$\boldsymbol{\tau}_1$ is recognizable; and*
(iii) *If $\boldsymbol{\tau}$ is M-quasi-invertible, then $\boldsymbol{\tau}_1$ is M-quasi-invertible. Conversely, if $\boldsymbol{\tau}$ is quasi-invertible, and $\boldsymbol{\tau}_1$ is M-quasi-invertible, then $\boldsymbol{\tau}$ is M-quasi-invertible.*

Proof.

(1) Note that if $1 \in R$ corresponds to $\boldsymbol{w}$ in $\mathcal{R}$, then $\boldsymbol{w}u_0$ is a prefix of $\boldsymbol{u}$, and so is $\boldsymbol{\tau}^n(u_0)$ for each n. Choose n large enough so that $|\boldsymbol{\tau}^n(u_0)| > |\boldsymbol{w}|$, so that $\boldsymbol{w}u_0$ is a prefix of $\boldsymbol{\tau}^n(u_0)$. If $j \in R$, and $\boldsymbol{w}' \in \mathcal{R}$ corresponds to j, then $\boldsymbol{w}'$ begins with u_0, so that $\boldsymbol{\tau}^n(u_0)$ and so $\boldsymbol{w}$ is a prefix of $\boldsymbol{\tau}^n(\boldsymbol{w}')$. Hence 1 is the first letter of $\boldsymbol{\tau}_1(j)$.
Since $\psi \circ \boldsymbol{\tau}_1 = \boldsymbol{\tau} \circ \psi$, then $\boldsymbol{\tau}_1(D(\boldsymbol{u})) = D(\boldsymbol{u}))$. As $\boldsymbol{\tau}_1$ is left proper, then it is primitive. If $\boldsymbol{\tau}_1(i) = \boldsymbol{\tau}_1(j)$, then either $\boldsymbol{\tau}(\alpha) = \boldsymbol{\tau}(\beta)$ for some $\alpha \neq \beta$, contradicting injectivity of $\boldsymbol{\tau}$, or the conditions in Theorem 3 are satisfied, contradicting recognizability of $\boldsymbol{\tau}$.

(2) Suppose that $\boldsymbol{\tau}_1$ is not recognizable. Then by Theorem 3, there exist letters i and j in R and some $\boldsymbol{y} \in Y$ with $\boldsymbol{\tau}_1(i)$ a proper suffix of $\boldsymbol{\tau}_1(j)$, and $\boldsymbol{\tau}_1(i)\boldsymbol{y}$ and $\boldsymbol{\tau}_1(j)\boldsymbol{y}$ appearing in Y with the same 1-cutting of $\boldsymbol{y}$. Suppose that $\psi(i) = \boldsymbol{w} = w_1 \ldots w_l$ and

$\psi(j) = \boldsymbol{w}' = w'_1 \dots w'_l$. Find the smallest k in $\{0 \dots l-1\}$ such that $w_{l-k} = w'_{l'-k}$. Such a k exists since otherwise $\boldsymbol{w}$ is a suffix of $\boldsymbol{w}'$, and since $\boldsymbol{w} \neq \boldsymbol{w}'$, then $w_1 = u_0$ occurs in $w'_2 \dots w'_l$, a contradiction to Part 2 of the definition of a return word. Now as $\boldsymbol{\tau}_1(i)$ is a proper suffix of $\boldsymbol{\tau}_1(j)$, there exist words $\boldsymbol{v}$ and $\boldsymbol{v}'$ in R^+, with $\boldsymbol{v}$ a suffix of $\boldsymbol{v}'$, $\boldsymbol{v}$ a suffix of $\boldsymbol{\tau}_1(i)$, $\boldsymbol{v}'$ a suffix of $\boldsymbol{\tau}_1(j)$, and words $\boldsymbol{p}$, $\boldsymbol{p}'$ and $\boldsymbol{s} \in \mathcal{A}^+$, with $\psi(\boldsymbol{v}) = \boldsymbol{p}\boldsymbol{\tau}(w_{l-k})\boldsymbol{s}$ and $\psi(\boldsymbol{v}') = \boldsymbol{p}'\boldsymbol{\tau}(w'_{l'-k})\boldsymbol{s}$. If $|\boldsymbol{\tau}(w_{l-k})| < |\boldsymbol{\tau}(w'_{l'-k})|$, then $\boldsymbol{\tau}(w_{l-k})$ is a proper suffix of $\boldsymbol{\tau}(w'_{l'-k})$; and if $|\boldsymbol{\tau}(w_{l-k})| > |\boldsymbol{\tau}(w'_{l'-k})|$, then $\boldsymbol{\tau}(w'_{l'-k})$ is a proper suffix of $\boldsymbol{\tau}(w_{l-k})$. Finally if $|\boldsymbol{\tau}(w_{l-k})| = |\boldsymbol{\tau}(w'_{l'-k})|$, then $\boldsymbol{\tau}(w_{l-k}) = \boldsymbol{\tau}(w'_{l'-k})$, which is not possible as $\boldsymbol{\tau}$ is injective on letters.

Thus (wlog) $\boldsymbol{\tau}(w_{l-k})$ is a proper subword of $\boldsymbol{\tau}(w'_{l'-k})$, and if $\boldsymbol{y}' = \boldsymbol{\tau}(w_{l-k+1} \dots w_l)\psi(\boldsymbol{y})$, then $\boldsymbol{\tau}(w_{l-k})\boldsymbol{y}'$ and $\boldsymbol{\tau}(w'_{l'-k})\boldsymbol{y}'$ appear in X with the same 1-cutting, a contradiction to the recognizability of $\boldsymbol{\tau}$.

(3) Note that if $\boldsymbol{x} \in X_{\boldsymbol{\tau}}$ has distinct σ-preimages $\boldsymbol{x}_1, \boldsymbol{x}_2, \dots \boldsymbol{x}_M$, then in $[u_0]$, there exist $\boldsymbol{y}_1, \boldsymbol{y}_2, \dots \boldsymbol{y}_M$ with $\boldsymbol{y}_i = \sigma^{-t_i}\boldsymbol{x}_i$, where $-t_i$ is the first time a preimage of $\boldsymbol{x}_i$ is in $[u_0]$, - that this point is well-defined follows from the quasi-invertibility and the aperiodicity of $(X_{\boldsymbol{\tau}}, \sigma)$. Let $\boldsymbol{y} = \sigma_{[u_0]}(\boldsymbol{x})$ if $\boldsymbol{x} \notin [u_0]$, and $\boldsymbol{x} = \boldsymbol{y}$ otherwise. Then $\boldsymbol{y}$ has M distinct $\sigma_{[u_0]}$-preimages, $\boldsymbol{y}_1, \dots, \boldsymbol{y}_M$. Since $\boldsymbol{x}$ is unique, so is $\boldsymbol{y}$. Now use the isomorphism in Lemma 17 to transfer this information to (Y, σ).

Conversely, if $(X_{\boldsymbol{\tau}}, \sigma)$ is N-quasi-invertible and $\boldsymbol{x}$ has N σ-preimages, $\{\boldsymbol{x}_1, \dots \boldsymbol{x}_N\}$, then if t is the first re-entry time of $\boldsymbol{x}$ into $[u_0]$, $\sigma^t(\boldsymbol{x})$ has N $\sigma_{[u_0]}$-preimages $\{\sigma^{-t_1}(\boldsymbol{x}_1), \dots \sigma^{-t_N}\boldsymbol{x}_N\}$. Thus $\psi^{-1}(\sigma^t(\boldsymbol{x}))$ has N σ-preimages, and as $(X_{\boldsymbol{\tau}_1}, \sigma)$ is M-quasi-invertible, $N = M$. ■

THEOREM 19.

Suppose that $\boldsymbol{\tau}$ is M-quasi-invertible and reasonable, with a generating fixed point $\boldsymbol{u}$, and branch point $\boldsymbol{y}$.

(i) *If $y_0 = u_0$, then there exists a semi-proper, stationary Bratteli diagram $\mathcal{B}$ such that $(X_{\mathcal{B}}, V_{\mathcal{B}})$ is topologically conjugate to (X, σ).*

(ii) *If $y_0 \neq u_0$, then there exists a semi-proper, stationary Bratteli diagram $\mathcal{B}$ such that a quotient of $(X_{\mathcal{B}}, V_{\mathcal{B}})$, is topologically conjugate to (X, σ).*

Proof. Using the fixed point $\boldsymbol{u}$, and working with the induced system $([u_0], \sigma_{[u_0]})$, we use Lemma 18 to find $\boldsymbol{\tau}_1$ left proper, minimal, recognizable, injective and aperiodic, such that $([u_0], \sigma_{[u_0]})$ is conjugate to $(X_{\boldsymbol{\tau}_1}, \sigma)$. If $\boldsymbol{y} \in [u_0]$, $(X_{\boldsymbol{\tau}}, \sigma)$ is a primitive of $([u_0], \sigma_{[u_0]})$. By Part 3 of Lemma 18, $\boldsymbol{\tau}_1$ is M-quasi-invertible. Now using Corollary 11, there is some $\boldsymbol{\tau}^*$ left proper, reasonable and quasi-invertible, with its fixed point $\mathbf{u}^*$ as branch point, such that $(X_{\boldsymbol{\tau}_1}, \sigma) = (X_{\boldsymbol{\tau}^*}, \sigma)$. If $\mathcal{B}^*$ is the semi-proper ordered Bratteli diagram associated to $\boldsymbol{\tau}^*$, then $(X_{\boldsymbol{\tau}^*}, \sigma)$, and so $(X_{\boldsymbol{\tau}_1}, \sigma)$, is isomorphic to $(X_{\mathcal{B}^*}, V_{\mathcal{B}^*})$. Using Theorem 15, $(X_{\boldsymbol{\tau}}, \sigma)$ is isomorphic to $(X_{\mathcal{B}'}, V_{\mathcal{B}'})$ where $\mathcal{B}'$ is obtained from $\mathcal{B}^*$ by the addition of a finite number of edges to $\mathcal{E}_1$.

If $\boldsymbol{y} \notin [u_0]$, then using Lemma 16, $(X_{\boldsymbol{\tau}}, \sigma)$ is a quotient of the primitive (X', σ) of $(X_{\boldsymbol{\tau}^*}, \sigma)$, and $(X_{\mathcal{B}'}, V_{\mathcal{B}'})$ is isomorphic to (X', σ). The result follows. ■

What properties do these ‘pinched’ adic systems of Part 2 of Theorem 19 have? It seems that even if superficially $\boldsymbol{\tau}$ has a pinched adic representation, it may still have an adic representation. If the branch point $\boldsymbol{y}$ satisfies $\mathrm{s}_1\boldsymbol{\tau}(\boldsymbol{y}) = \boldsymbol{y}$, then is $\boldsymbol{y}$ always the fixed point of some $\boldsymbol{\tau}^*$, which has some more general definition than the one in Lemma 10? We illustrate with Example (3) below. Also, do these results extend to substitutions with several branch points, as in Example (4) below?

Examples:

(1) If $\boldsymbol{\tau}$ is any reasonable suffix permutative substitution, with one fixed point $\boldsymbol{u}$, then it is quasi-invertible, and $(X_{\boldsymbol{\tau}}, \sigma)$ is a primitive over $(X_{\boldsymbol{\tau}_1}, \sigma)$, as $\phi(\boldsymbol{u}) \in X_{\boldsymbol{\tau}_1}$.

(2) If $\boldsymbol{\tau}(a) = aac$, $\boldsymbol{\tau}(b) = bcc$, and $\boldsymbol{\tau}(c) = abc$, then $(X_{\boldsymbol{\tau}}, \sigma)$ is 3-quasi-invertible, with the non-invertible element $\boldsymbol{y}$ satisfying $c\boldsymbol{\tau}(y) = y$. The two $\boldsymbol{\tau}$-fixed points $\boldsymbol{u}$ and $\boldsymbol{v}$ are in $[a] \cup [b]$; so if $U = [a]$ (or $[b]$), $(X_{\boldsymbol{\tau}}, \sigma)$ is recoverable only as a quotient of a primitive of (U, σ_U). However Lemma 10 applies, since all $\boldsymbol{\tau}$-words have a proper common suffix. Thus Corollary 11 tells us that $(X_{\boldsymbol{\tau}}, \sigma) = (X_{\boldsymbol{\tau}^*}, \sigma)$ where $\boldsymbol{\tau}^*(\boldsymbol{y}) = \boldsymbol{y}$, now $(X_{\boldsymbol{\tau}^*}, \sigma)$ is a primitive over $[c]$.

(3) If $\boldsymbol{\tau}(a) = aac$, $\boldsymbol{\tau}(b) = bcc$, $\boldsymbol{\tau}(c) = adbc$ and $\boldsymbol{\tau}(d) = adbd$, then since $da \notin \mathcal{L}_{\boldsymbol{\tau}}$ and $cb \notin \mathcal{L}_{\boldsymbol{\tau}}$, both fixed points are σ-invertible. Inspection of maximal proper common suffixes of the $\boldsymbol{\tau}$-substitution words of letters in any subset $\mathcal{A}^*$ of $\mathcal{A}$ leads to the existence of only one branch point $\boldsymbol{y}$ satisfying $c\boldsymbol{\tau}(\boldsymbol{y}) = \boldsymbol{y}$. Lemma 10 cannot directly apply because the family of $\boldsymbol{\tau}$-substitution words does not have a common suffix.
If we consider instead the induced system $([c], \sigma_{[c]})$, $(X_{\boldsymbol{\tau}}, \sigma)$ is a primitive over $([c], \sigma_{[c]})$. Define $\boldsymbol{\tau}^*(a) = caa$, $\boldsymbol{\tau}^*(b) = c$, $\boldsymbol{\tau}^*(c) = cadb$ and $\boldsymbol{\tau}^*(d) = cadbdb$. One can now prove, similarly to the proof of Lemma 10 to show that $\boldsymbol{\tau}^*(\boldsymbol{y}) = \boldsymbol{y}$. Thus $([c], \sigma_{[c]})$ is itself a (left proper, quasi-invertible, reasonable) substitution subshift, and so has a stationary adic representation $(X_{\mathcal{B}^*}, V_{\mathcal{B}^*})$. Now Theorem 15 applies, so that $(X_{\boldsymbol{\tau}}, \sigma)$ also has a stationary adic representation. In this example it is relatively straightforward to find the right $\boldsymbol{\tau}^*$; in general though it is not clear how a branch point $\boldsymbol{y}$ can be seen as the fixed point of some $\boldsymbol{\tau}^*$. If this were the case, then all quasi-invertible, aperiodic, minimal, injective substitutions would have an adic representation.

(4) If $\boldsymbol{\tau}(0) = 01$ and $\boldsymbol{\tau}(1) = 10$ (the *Morse* substitution), then there are two $\boldsymbol{\tau}$-fixed points $\boldsymbol{u} \in [0]$ and $\overline{\boldsymbol{u}} \in [1]$, and these are the only branch points. The set of return words to 0 in $\boldsymbol{u}$ are $\mathcal{R} = \{011, 01, 0\}$, and $\boldsymbol{\tau}_1^2$ is defined by $\boldsymbol{\tau}_1^2(1) = 123132$, $\boldsymbol{\tau}_1^2(2) = 1232$ and $\boldsymbol{\tau}_1^2(3) = 13$, and $\boldsymbol{\tau}_1^2$ has 2 branch points, the $\boldsymbol{\tau}_1$-fixed point, and a point $\boldsymbol{y}$ satisfying $32\boldsymbol{\tau}(\boldsymbol{y}) = \boldsymbol{y}$. If we take $\mathcal{R}$ to be the set of return words to 1 in $\overline{\boldsymbol{u}}$, then the resulting $\boldsymbol{\tau}_1$ is the same. Neither $\boldsymbol{\tau}$ nor $\boldsymbol{\tau}_1$ currently has an appropriate adic representation, so this technique fails here.

(5) *Minimal rank one subshifts* are defined by substitutions on $\mathcal{A} = \{0, 1\}$ of the form

$$\boldsymbol{\tau}(0) = 0^{n_1} 1^{m_1} 0^{n_2} 1^{m_2} \ldots 0^{n_{k-1}} 1^{m_{k-1}} 0^{n_k} \text{ and } \boldsymbol{\tau}(1) = 1,$$

where $k < \infty$, and n_1 and n_k are positive. The latter condition ensures that the resulting substitution subshift is minimal, with generating fixed point $\boldsymbol{u} := \lim_{n\to\infty} \boldsymbol{\tau}^n(0)$. These systems are equivalent to rank-one systems defined by 'cutting and stacking' where there are a bounded number of cuts and spacers added, and the same cutting-and-stacking rule is obeyed at each stage. A comprehensive exposition of rank one systems is given in [**Fer97**]. Suppose $n_i \neq n_j$ for some i, j, so that $\boldsymbol{\tau}$ is aperiodic. Then $(X_{\boldsymbol{\tau}}, \sigma)$ is quasi-invertible if and only if $m_1 = m_2 = \ldots = m_{k-1} = m > 0$. For, the generating fixed point $\boldsymbol{u}$ is a branch point, and the fixed point $1^{m_i}\boldsymbol{u}$ is another branch point if and only $m_i < m_j$ for some j.
If $\boldsymbol{\tau}(0) = 0^{n_1} 1^m 0^{n_2} 1^m \ldots 0^{n_{k-1}} 1^m 0^{n_k}$, then there are two return words to 0, $\boldsymbol{w}_1 = 0$ and $\boldsymbol{w}_2 = 01^m$. In this case $\boldsymbol{\tau}_1$ is defined on $\{1, 2\}$ and $\boldsymbol{\tau}_i = \boldsymbol{w}i$ where $\boldsymbol{w} = 1^{n_1-1} 2 1^{n_2-1} 2 \ldots 1^{n_k-1}$. By Lemma 18, $\boldsymbol{\tau}_1$ is left proper, reasonable and 2-quasi-invertible. Thus $(X_{\boldsymbol{\tau}}, \sigma)$ is a primitive of $([0], \sigma_{[0]})$ and by Theorem 19, $\boldsymbol{\tau}$ has a semi proper adic representation by adding $m - 1$ edges from v_0 to the vertex in $\mathcal{V}_1$ corresponding to the letter '1' in the Bratteli representation of $\boldsymbol{\tau}_1$. For $1 \leq k \leq M$, the $\boldsymbol{\tau}$-fixed points $1^k\boldsymbol{u}$ live on (different) levels of the tower over $(X_{\boldsymbol{\tau}}, \sigma)$.
If for some i, j, $m_i \neq m_j$, then $\boldsymbol{\tau}_1$ is still of the form $\boldsymbol{\tau}_1(i) = \boldsymbol{w}i$, where $\boldsymbol{w} = 1^{n_1-1} 2 1^{n_2-1} 3 \ldots 1^{n_{k-1}-1} k 1^{n_k-1}$, so that $\boldsymbol{\tau}_1$ is M-quasi-invertible, with $M \geq 3$. Thus $(X_{\boldsymbol{\tau}_1}, \sigma)$ is quasi-invertible, even though $(X_{\boldsymbol{\tau}}, \sigma)$ is not, and we cannot generate an

adic representation for $\boldsymbol{\tau}$ from one for $\boldsymbol{\tau}_1$. For example, if $\boldsymbol{\tau}(0) = 00100110$, then the return words are $\mathcal{R} = \{0, 01, 011\}$ and $\boldsymbol{\tau}_1$ is 3-almost periodic, with $\boldsymbol{x}^1, \boldsymbol{x}^2$, and $\boldsymbol{x}^3$ the preimages of $\mathcal{D}(\boldsymbol{u})$. In X though, $\phi(\boldsymbol{x}^1)$, and $\sigma(\phi(\boldsymbol{x}^2))$ are preimages of $\boldsymbol{u}$, and $\phi(\boldsymbol{x}^2)$ and $\sigma(\phi(\boldsymbol{x}^3))$ are preimages of $\sigma(\phi(\boldsymbol{x}^2))$. So an adic representation of $(X_{\boldsymbol{\tau}}, \sigma)$ would need 4 maximal elements, one of which is also minimal, and another minimal element.

References

AR91 Pierre Arnoux and Gérard Rauzy. Représentation géométrique de suites de complexité $2n+1$. *Bull. Soc. Math. France*, 119(2):199–215, 1991.

BKM09 S. Bezuglyi, J. Kwiatkowski, and K. Medynets. Aperiodic substitution systems and their Bratteli diagrams. *Ergodic Theory Dynam. Systems*, 29(1):37–72, 2009.

CDHM03 Maria Isabel Cortez, Fabien Durand, Bernard Host, and Alejandro Maass. Continuous and measurable eigenfunctions of linearly recurrent dynamical Cantor systems. *J. London Math. Soc. (2)*, 67(3):790–804, 2003.

CMPS93 D. Crisp, W. Moran, A. Pollington, and P. Shiue. Substitution invariant cutting sequences. *J. Théor. Nombres Bordeaux*, 5(1):123–137, 1993.

DHS99 F. Durand, B. Host, and C. Skau. Substitutional dynamical systems, Bratteli diagrams and dimension groups. *Ergodic Theory Dynam. Systems*, 19(4):953–993, 1999.

Fer97 Sébastien Ferenczi. Systems of finite rank. *Colloq. Math.*, 73(1):35–65, 1997.

For97 A. H. Forrest. K-groups associated with substitution minimal systems. *Israel J. Math.*, 98:101–139, 1997.

GJ00 Richard Gjerde and Ørjan Johansen. Bratteli-Vershik models for Cantor minimal systems: applications to Toeplitz flows. *Ergodic Theory Dynam. Systems*, 20(6):1687–1710, 2000.

GJ02 Richard Gjerde and Ørjan Johansen. Bratteli-Vershik models for Cantor minimal systems associated to interval exchange transformations. *Math. Scand.*, 90(1):87–100, 2002.

GPS95 Thierry Giordano, Ian F. Putnam, and Christian F. Skau. Topological orbit equivalence and C^*-crossed products. *J. Reine Angew. Math.*, 469:51–111, 1995.

Hos86 B. Host. Valeurs propres des systèmes dynamiques définis par des substitutions de longueur variable. *Ergodic Theory Dynam. Systems*, 6(4):529–540, 1986.

HPS92 Richard H. Herman, Ian F. Putnam, and Christian F. Skau. Ordered Bratteli diagrams, dimension groups and topological dynamics. *Internat. J. Math.*, 3(6):827–864, 1992.

Mos92 Brigitte Mossé. Puissances de mots et reconnaissabilité des points fixes d'une substitution. *Theoret. Comput. Sci.*, 99(2):327–334, 1992.

PS73 Karl Petersen and Leonard Shapiro. Induced flows. *Trans. Amer. Math. Soc.*, 177:375–390, 1973.

Que87 Martine Queffélec. *Substitution dynamical systems—spectral analysis*, volume 1294 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1987.

Sen06 E. Seneta. *Non-negative matrices and Markov chains*. Springer Series in Statistics. Springer, New York, 2006. Revised reprint of the second (1981) edition [Springer-Verlag, New York; MR0719544].

Ver81 A. M. Vershik. Uniform algebraic approximation of shift and multiplication operators. *Dokl. Akad. Nauk SSSR*, 259(3):526–529, 1981.

VL92 A. M. Vershik and A. N. Livshits. Adic models of ergodic transformations, spectral theory, substitutions, and related topics. In *Representation theory and dynamical systems*, volume 9 of *Adv. Soviet Math.*, pages 185–204. Amer. Math. Soc., Providence, RI, 1992.

Dept. of Mathematics, Trent University
1600 West Bank Drive, Peterborough,
Ontario, K9J 7B8 Canada.

ryassawi@trentu.ca